

# ON THE OPERATOR EQUATION $AX - XB = C$ WITH UNBOUNDED OPERATORS $A$ , $B$ , AND $C$

NGUYEN THANH LAN

Received 30 July 2001

We find the criteria for the solvability of the operator equation  $AX - XB = C$ , where  $A$ ,  $B$ , and  $C$  are unbounded operators, and use the result to show existence and regularity of solutions of nonhomogeneous Cauchy problems.

## 1. Introduction

Let  $A$  and  $B$  be operators on Banach spaces  $E$  and  $F$ , respectively, and let  $C$  be an operator from  $F$  to  $E$ . Of concern is the operator equation

$$AX - XB = C. \quad (1.1)$$

To be found is a bounded operator  $X$  from  $F$  to  $E$  such that  $X(D(B) \cap D(C)) \subseteq D(A)$  and  $AXf - XBf = Cf$  for every  $f \in D(B) \cap D(C)$ . Over the last few decades, (1.1) has been considered by many authors. It was first studied intensively for bounded operators by Daleckii and Krein [2], Rosenblum [16] (see also [5]). For unbounded operators  $A$  and  $B$ , the case when  $A$  and  $B$  are generators of  $C_0$ -semigroups was considered in [1, 4, 10]. Recently, many papers apply the results to the stability and regularity of solutions of the abstract differential equation

$$u'(t) = Au(t) + f(t), \quad (1.2)$$

(see [10, 12, 13]), and the higher differential equation

$$u^{(n)}(t) = Au(t) + f(t) \quad (1.3)$$

(see [8, 17]). On the other hand, it seems that there is little consideration of (1.1) when  $C$  is an unbounded operator.

In this paper, we study (1.1) for this case. The motivation behind this is that, if  $X$  is a bounded solution of (1.1), then the operator  $\mathcal{A}' := \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$  is similar to the operator  $\mathcal{A} := \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$  by the identity

$$\begin{pmatrix} A & C \\ 0 & B \end{pmatrix} = \begin{pmatrix} I & X \\ 0 & I \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \begin{pmatrix} I & -X \\ 0 & I \end{pmatrix}. \quad (1.4)$$

Therefore, many properties of  $\mathcal{A}$  can be carried over to  $\mathcal{A}'$ . Thus, instead of studying the unbounded perturbed operator  $\mathcal{A}'$  we just study the operator  $\mathcal{A}$ , which seems to be much simpler. In particular, the operator  $\mathcal{A}'$  is a generator of a  $C_0$ -semigroup on  $E \times F$  if and only if  $\mathcal{A}$  is a generator of a  $C_0$ -semigroup on  $E \times F$ . It is very useful, when converting the nonhomogeneous Cauchy problem

$$u'(t) = Au(t) + f(t), \quad t \geq 0, \quad u(0) = x_0 \in E, \quad (1.5)$$

where  $f$  is a vector of a function space  $F(\mathbb{R}, E)$ , into a homogeneous problem

$$\mathcal{U}'(t) = \begin{pmatrix} A & \delta_0 \\ 0 & \frac{d}{dt} \end{pmatrix} \mathcal{U}(t), \quad t \geq 0, \quad \mathcal{U}(0) = (x_0, f). \quad (1.6)$$

on  $E \times F(\mathbb{R}, E)$  (see [6, 7, 9]). Note that the operator  $\delta_0$  is unbounded in some function space  $F(\mathbb{R}, E)$ .

We organize this paper as follows: in Section 2, we first show the solvability of (1.1). Then we give some applications to the existence and regularity of solutions of the nonhomogeneous Cauchy problem. In Section 3, we consider the nonhomogeneous differential equation

$$u'(t) = Au(t) + f(t) \quad (1.7)$$

on the line  $\mathbb{R}$ , where  $f \in L^p(\mathbb{R}, E)$ . It turns out that the existence and uniqueness of the bounded mild solution of (1.7) is equivalent to the solvability of equation  $AX - XB = \delta_0$ . (See the notations below.)

We fix some notations. Let  $E$  be a Banach space. The value of a functional  $\phi \in E^*$  at a vector  $x \in E$  is denoted by  $\langle x, \phi \rangle$ . By  $W^{p,1}(\mathbb{R}, E)$  we denote the space of all absolutely continuous functions  $f$  from  $\mathbb{R}$  to  $E$  with  $f' \in L^p(\mathbb{R}, E)$ . If  $F(\mathbb{R}, E)$  is a certain function space over  $E$ , then  $\mathcal{D} : D(\mathcal{D}) \subseteq F(\mathbb{R}, E) \rightarrow F(\mathbb{R}, E)$  is defined by  $\mathcal{D}f = f'$  and  $\delta_0 : D(\delta_0) \subseteq F(\mathbb{R}, E) \rightarrow E$  by  $\delta_0(f) = f(0)$ . Finally, for  $\lambda \in \varrho(A)$ ,  $(\lambda - A)^{-1}$  is denoted by  $R(\lambda, A)$ .

## 2. Solution of the equation $AX - XB = C$

Throughout this paper,  $A$  and  $(-B)$  will denote generators of  $C_0$ -semigroups  $(T(t))$  and  $(S(t))$  on Banach spaces  $E$  and  $F$ , respectively, and  $C$  is an operator from  $F$  to  $E$ . For the operator equation

$$AX - XB = C, \quad (2.1)$$

we recall a definition. Let  $B$  be a linear operator on  $F$ . Then we say that  $C : F \rightarrow E$  is *B-bounded* if  $D(B) \subseteq D(C)$  and the operator  $C(\lambda - B)^{-1}$  is bounded for one (all)  $\lambda \in \varrho(B)$ . For example, if  $C$  is a closed operator with  $D(B) \subseteq D(C)$ , then  $C$  is *B-bounded*. We have the following theorem about the solvability of (2.1).

**THEOREM 2.1.** *Let  $A$  and  $-B$  be generators of  $C_0$ -semigroups  $(T(t))_{t \geq 0}$  and  $(S(t))_{t \geq 0}$  on  $E$  and  $F$ , respectively, and  $C$  *B-bounded*. Let*

$$\begin{aligned} Q(t) : F &\supseteq D(B) \mapsto E : Q(t)f := T(t)CS(t)f, \quad t \geq 0, \\ R(t) : F &\supseteq D(B) \mapsto E : R(t)f := - \int_0^t Q(s)f ds, \quad t \geq 0. \end{aligned} \quad (2.2)$$

We assume that

- (i) the weak-topology closure of  $\{Q(t)f\}_{t \geq 0}$  contains 0 for every  $f \in D(B)$ ;
- (ii)  $R(t)$  can be extended to a bounded operator for every  $t \geq 0$  and the family  $\{R(t)\}_{t \geq 0}$  is relatively compact in the weak operator topology.

Then (2.1) has a bounded solution.

Conversely, if (2.1) has a bounded solution, then  $R(t)$  is bounded for every  $t \geq 0$ . In addition, if for every bounded operator  $Y$  from  $F$  to  $E$ ,  $T(t)YS(t)$  converges to 0 as  $t \rightarrow \infty$  in the weak (strong, uniform) operator topology, then the solution  $X$  of (2.1) is unique, and  $R(t)$  converges to  $X$  in the weak (strong, uniform) operator topology.

**Remark 2.2.** The operator  $R(t)$  is meaningful for each  $t \geq 0$ , since the function  $t \rightarrow Q(t)f = T(t)CS(t)f = T(t) \cdot CR(\lambda, B) \cdot S(t) \cdot (\lambda - B)f$  is continuous for each  $f \in D(B)$  and  $\lambda \in \varrho(B)$ .

*Proof.* Let  $\lambda \in \varrho(B)$  and take  $C_1 = CR(\lambda, B)$ . Then  $C_1$  is bounded. For  $t \geq 0$ , we define the operators  $Q_1(t) : F \rightarrow E$  and  $R_1(t) : F \rightarrow E$  by

$$\begin{aligned} Q_1(t)f &:= T(t)C_1S(t)f, \\ R_1(t)f &:= - \int_0^t Q_1(s)f ds. \end{aligned} \quad (2.3)$$

Then  $Q_1(t)$  and  $R_1(t)$  are bounded operators. We now consider the operator equation

$$AY - YB = C_1. \quad (2.4)$$

By assumptions, there exists a net  $t_\alpha \rightarrow \infty$  such that  $T(t_\alpha)CS(t_\alpha)$  converges weakly to 0 and  $R(t_\alpha)$  converges weakly to a bounded operator  $Q$ . Therefore,  $T(t_\alpha)C_1S(t_\alpha)$  converges weakly to 0 and  $R_1(t_\alpha)$  converges weakly to the bounded operator  $QR(\lambda, B)$ . By [10, Theorem 3], (2.4) has a bounded solution, namely  $Y = QR(\lambda, B)$ . It implies that  $Y(\lambda - B)$  can be extended to the bounded operator  $Q$ . We verify that  $Q = \overline{Y(\lambda - B)}$  is a solution of (2.1).

First, for any  $f \in D(B^2)$  we have  $(\lambda - B)f \in D(B)$  and

$$\begin{aligned} A\mathcal{Q}f - QBf &= AY(\lambda - B)f - Y(\lambda - B)Bf \\ &= (AY - YB)(\lambda - B)f \\ &= C_1(\lambda - B)f = Cf. \end{aligned} \quad (2.5)$$

Hence,  $A\mathcal{Q}f - QBf = Cf$  and thus,

$$T(t)A\mathcal{Q}S(t)f - T(t)QS(t)Bf = T(t)CS(t)f \quad (2.6)$$

for all  $f \in D(B^2)$ . By [10, Lemma 1], if  $f \in D(B^2)$  and  $\phi \in D(A')$  we have

$$\begin{aligned} \frac{d}{dt} \langle T(t)QS(t)f, \phi \rangle &= \langle T(t)A\mathcal{Q}S(t)f, \phi \rangle - \langle T(t)QS(t)Bf, \phi \rangle \\ &= \langle T(t)CS(t)f, \phi \rangle. \end{aligned} \quad (2.7)$$

Therefore,

$$\begin{aligned} \langle R(t)f, \phi \rangle &= - \int_0^t \langle T(s)CS(s)f, \phi \rangle ds \\ &= - \int_0^t \frac{d}{ds} \langle T(s)QS(s)f, \phi \rangle ds \\ &= - \langle T(t)XS(t)f - Qf, \phi \rangle, \end{aligned} \quad (2.8)$$

from which it follows that

$$R(t)f = Qf - T(t)QS(t)f \quad (2.9)$$

for  $f \in D(B^2)$ . Since the operators on both sides of (2.9) are bounded and  $D(B^2)$  is dense in  $F$ , it implies that (2.9) also holds for all  $f \in D(B)$ .

Let now  $f \in D(B)$  and  $\phi \in D(A')$ , then we have

$$\begin{aligned} \langle T(t)CS(t)f, \phi \rangle &= \frac{d}{dt} \int_0^t \langle T(s)CS(s)f, \phi \rangle ds = - \frac{d}{dt} \langle R(t)f, \phi \rangle \\ &= \frac{d}{dt} \langle T(t)QS(t)f - Qf, \phi \rangle = \frac{d}{dt} \langle T(t)QS(t)f, \phi \rangle \\ &= \langle T(s)A\mathcal{Q}S(s)f - T(s)QS(s)Bf, \phi \rangle, \end{aligned} \quad (2.10)$$

which implies

$$T(t)A\mathcal{Q}S(t)f - T(t)QS(t)Bf = T(t)CS(t)f \quad \forall t \geq 0. \quad (2.11)$$

Taking  $t = 0$  we have  $A\mathcal{Q}f - QBf = Cf$  for  $f \in D(B)$ .

Conversely, if  $X$  is a solution of (2.1), then by the same argument as above we have

$$R(t)f = Xf - T(t)XS(t)f \quad (2.12)$$

for  $f \in D(B)$ . Since all the operators on the right-hand side of (2.12) are bounded and  $D(B)$  is dense in  $F$ ,  $R(t)$  can be extended to a bounded operator. Moreover, if  $T(t)XS(t) \rightarrow 0$  in weak (resp., strong, uniform) operator topology, then  $R(t) \rightarrow X$  weakly (resp., strongly, uniformly). Hence,  $X$  is uniquely determined, and the proof is complete.  $\square$

For a semigroup  $(T(t))_{t \geq 0}$  generated by  $A$ , we define *the growth bound*  $\omega(A)$  by

$$\omega(A) := \inf \{ \lambda \in \mathbb{R} : \exists M > 0 \text{ such that } \|T(t)\| \leq M e^{\lambda t} \forall t \geq 0 \}. \quad (2.13)$$

If  $\omega(A) < 0$ , then  $(T(t))$  is called *uniformly exponentially stable*. The following is a short version of [Theorem 2.1](#), which gives the existence and uniqueness of the solution of (2.1) and will be used more frequently.

**THEOREM 2.3.** *Assume that  $\omega(A) + \omega(-B) < 0$  and that  $R(t)$  is uniformly bounded. Then (2.1) has a unique bounded solution.*

*Proof.* Since  $AX - XB = (A + \lambda)X - X(B + \lambda)$ , we can assume, without loss of generality, that  $\omega(S) = 0$  and  $\omega(T) < 0$ . Then for any  $\lambda \in \varrho(B)$  we have

$$\begin{aligned} \|T(t)CS(t)f\| &= \|T(t)CR(\lambda, B)S(t)(\lambda - B)f\| \\ &\leq M_1 e^{\omega(A)t} \cdot \|CR(\lambda, B)\| \cdot M_2 \cdot \|(\lambda - B)f\| \longrightarrow 0. \end{aligned} \quad (2.14)$$

So [Theorem 2.1](#)(i) is satisfied. In addition, for  $t_1, t_2 \rightarrow \infty$  and  $f \in D(B)$  we have

$$\begin{aligned} \|R(t_1)f - R(t_2)f\| &\leq \int_{t_1}^{t_2} \|T(s)CS(s)f\| ds \\ &\leq \int_{t_1}^{t_2} M_1 e^{\omega(A)s} \|CR(\lambda, B)\| M_2 \|(\lambda - B)f\| ds \\ &\leq M \int_{t_1}^{t_2} e^{\omega(A)s} ds \longrightarrow 0 \quad \text{as } t_1, t_2 \longrightarrow \infty. \end{aligned} \quad (2.15)$$

Since  $R(t)$  is uniformly bounded and  $D(B)$  is dense in  $F$ ,  $R(t)$  converges strongly to a bounded operator. So [Theorem 2.1](#)(ii) is satisfied. By [Theorem 2.1](#), (1.1) has a solution, and since  $\|T(t)YS(t)f\| \rightarrow 0$ ,  $t \rightarrow \infty$  for each bounded operator  $Y : F \rightarrow E$ , it is unique and equals to the bounded extension of  $\lim_{t \rightarrow \infty} R(t)$ .  $\square$

The following corollary, which is involved with *exponentially dichotomic* semigroups follows directly from [Theorem 2.3](#). Recall, a  $C_0$ -semigroup  $(T(t))$  on a Banach space  $E$  is exponentially dichotomic, if there is a bounded projection  $P$  on  $E$  and positive constants,  $M$  and  $\omega$ , such that

- (i)  $PT(t) = T(t)P$  for all  $t \geq 0$ ;
- (ii)  $\|T(t)x\| \leq Me^{-\omega t}\|x\|$  for all  $x \in P(E)$ ;
- (iii) the restriction  $T(t)|_{\ker(P)}$  extends to a group and  $\|T(-t)x\| \leq Me^{-\omega t}\|x\|$  for all  $x \in \ker(P)$  and  $t \geq 0$ .

It is well known that  $(T(t))$  is exponentially dichotomic if and only if  $\sigma(T(t)) \cap \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \emptyset$  for all  $t > 0$  (cf. [14]). By the weak spectral mapping theorem, this implies that  $\sigma(A) \cap i\mathbb{R} = \emptyset$ . Also note that uniformly exponentially stable semigroup is a particular case of exponentially dichotomic one, when  $P = I$ , the identity operator.

**COROLLARY 2.4.** *Let  $A$  be the generator of an exponentially dichotomic semigroup  $(T(t))$  and  $-B$  the generator of an isometric  $C_0$ -group. If  $R(t)$  is uniformly bounded. Then (2.1) has a unique bounded solution.*

*Proof.* Let  $C_1 = PC$ ,  $C_2 = (I - P)C$ ,  $A_1 = A|_{P(E)}$ ,  $A_2 = A|_{\ker(P)}$ . It is easy to see that  $C_1$  and  $C_2$  are  $B$ -bounded. By [Theorem 2.3](#), there is a unique bounded operator  $X_1 : F \rightarrow P(E)$  such that

$$A_1 X_1 - X_1 B = C_1. \quad (2.16)$$

Moreover,  $-A_2$  generates an exponentially stable semigroup and  $B$  also generates an isometric group. Again, by [Theorem 2.3](#), there is a unique bounded operator  $X_2 : F \rightarrow \ker(P)$  such that

$$-A_2 X_2 + X_2 B = -C_2, \quad \text{or} \quad A_2 X_2 - X_2 B = C_2. \quad (2.17)$$

Let now  $X = X_1 + X_2$ , then  $AX - XB = A(X_1 + X_2) - (X_1 + X_2)B = (A_1 X_1 - X_1 B) + (A_2 X_2 - X_2 B) = C_1 + C_2 = C$ . Thus,  $X$  is a bounded solution of (2.1). The uniqueness of  $X$  follows from the fact that  $PX$  and  $(I - P)X$  are unique bounded solutions of (2.16) and (2.17), respectively.  $\square$

In the following we apply the above results to study the existence and regularity of solutions of Cauchy problems.

**COROLLARY 2.5.** *Let  $A$  be the generator of a  $C_0$ -semigroup  $(T(t))$  on  $E$  such that  $\omega(T) < 0$  and  $\mathcal{D} : W^{p,1}(\mathbb{R}, E) \rightarrow L^p(\mathbb{R}, E)$  be given by  $\mathcal{D} = d/dt$ . Then the equation*

$$AX - X\mathcal{D} = \delta_0 \quad (2.18)$$

*has a unique solution.*

*Proof.* It is well known that  $\delta_0$  is  $\mathcal{D}$ -bounded and  $-\mathcal{D}$  is the generator of the shift  $C_0$ -group  $S(t)$  given by  $S(t)f(s) = f(s - t)$ . Hence,  $\omega(S) = 0$ . Moreover,

$$\begin{aligned}
\|R(t)f\| &= \left\| \int_0^t T(s)\delta_0 S(s)f ds \right\| = \left\| \int_0^t T(s)f(-s) ds \right\| \\
&\leq M \int_0^t \|f(-s)\| ds \leq M \|f\|.
\end{aligned} \tag{2.19}$$

Hence  $R(t)$ ,  $t \geq 0$ , is uniformly bounded. Thus, by [Theorem 2.3](#), [\(2.18\)](#) has a unique bounded solution.  $\square$

From [Corollary 2.5](#) we obtain the following.

**COROLLARY 2.6.** *Let  $p \geq 1$  and  $A$  the generator of a  $C_0$ -semigroup in  $E$ . Then the operator*

$$\mathcal{A} := \begin{pmatrix} A & \delta_0 \\ 0 & \mathcal{D} \end{pmatrix} \tag{2.20}$$

*with  $D(\mathcal{D}) := D(A) \times W^{1,1}(\mathbb{R}, E)$  is the generator of a  $C_0$ -semigroup on  $E \times L^p(\mathbb{R}, E)$ .*

*Proof.* Without loss of generality, we assume  $\omega(T) < 0$ . Then, by [Corollary 2.5](#), [\(2.18\)](#) has a unique solution. Hence,  $\mathcal{A}$  is similar to the generator  $\begin{pmatrix} A & 0 \\ 0 & \mathcal{D} \end{pmatrix}$  and thus is a generator.  $\square$

It is well known (cf. [7]) that if  $(u_1, u_2)^T$  is a (classical) solution of the Cauchy problem

$$\mathcal{U}'(t) = \mathcal{A}\mathcal{U}(t) \quad t \geq 0, \quad \mathcal{U}(0) = (u_0, f)^T \tag{2.21}$$

on  $E \times F(\mathbb{R}, E)$ , where  $F(\mathbb{R}, E)$  is a function space, then the first component  $u_1$  is the (classical) solution of the inhomogeneous Cauchy problem

$$u'(t) = Au(t) + f(t) \quad t \geq 0, \quad u(0) = u_0. \tag{2.22}$$

From the above observation and [Corollary 2.6](#) we obtain the following.

**COROLLARY 2.7.** *Let  $A$  be the generator of a  $C_0$ -semigroup and  $f \in W^{1,1}(\mathbb{R}, E)$ , then [\(2.22\)](#) has a unique classical solution.*

Now recall the definition of *extrapolation space*. Let  $A$  be the generator of a  $C_0$ -semigroup  $(T(t))$  on a Banach space  $E$  and  $\lambda \in \varrho(A)$ . On  $E$  we introduce a new norm by

$$\|x\|_{-1} = \|R(\lambda, A)x\|. \tag{2.23}$$

Then the completion of  $(E, \|\cdot\|_{-1})$  is called the *extrapolation space* of  $E$  associated with  $A$ , and is denoted by  $E_{-1}$ . It is shown that the operator  $T(t)$  can

be uniquely extended to a bounded operator on the Banach space  $E_{-1}$ . The result is a  $C_0$ -semigroup on  $E_{-1}$ , denoted by  $(T_{-1}(t))$ . The semigroup  $(T_{-1}(t))$  is called the *extrapolated semigroup* of  $(T(t))$ . If we denote by  $A_{-1}$  the generator of  $(T_{-1}(t))$  on  $E_{-1}$ , then we have the following properties (see more details in [3, 7]):

- (i)  $\|T_{-1}(t)\|_{L(E_{-1})} = \|T(t)\|_{L(E)}$ ;
- (ii)  $E$  is dense in  $E_{-1}$  and  $D(A_{-1}) = E$ ;
- (iii)  $A_{-1} : E \rightarrow E_{-1}$  is the unique extension of  $A : D(A) \rightarrow E$  to  $E \rightarrow E_{-1}$ .

The following two corollaries show the existence and uniqueness of the classical solution of the nonhomogeneous Cauchy problem (2.22) for the case that the nonhomogeneous term  $f$  is not differentiable. Since their proofs are similar, we present here only one of them.

**COROLLARY 2.8.** *Let  $A$  be the generator of an analytic semigroup. Then (2.22) has a unique classical solution for every  $x \in D(A)$  and Hölder continuous function  $f \in H_\alpha(\mathbb{R}, E)$ .*

**COROLLARY 2.9.** *Let  $A$  be the generator of a  $C_0$ -semigroup on  $E$ . Then (2.22) has a unique classical solution for every  $x \in D(A)$  and  $f \in BUC(\mathbb{R}, [D(A)])$ , where  $[D(A)]$  is the Banach space  $(D(A), \|\cdot\|_A)$  with the norm  $\|x\|_A = \|x\| + \|Ax\|$ .*

*Proof of Corollary 2.9.* Without loss of generality, we assume  $\omega(A) < 0$ . In view of the previous observation we have only to show that  $\mathcal{A} := \begin{pmatrix} A & \delta_0 \\ 0 & \mathcal{D} \end{pmatrix}$  with  $D(\mathcal{A}) = D(A) \times BUC(\mathbb{R}, [D(A)])$  is a generator of a  $C_0$ -semigroup on  $E \times BUC_{-1}(\mathbb{R}, [D(A)])$ , where  $BUC_{-1}(\mathbb{R}, [D(A)])$  is the extrapolated space of  $BUC(\mathbb{R}, [D(A)])$  associated with  $\mathcal{D} = d/dt$  on  $BUC(\mathbb{R}, [D(A)])$ . This is done if we show that there is a bounded solution of the operator equation

$$AX - X\mathcal{D} = \delta_0, \quad (2.24)$$

where  $F := BUC_{-1}(\mathbb{R}, [D(A)])$ . It is easy to see that  $\delta_0$  is  $\mathcal{D}$ -bounded. Since  $\omega(\mathcal{D}) = 0$  we have  $\omega(A) + \omega(\mathcal{D}) < 0$ . Moreover, let  $f \in BUC(\mathbb{R}, [D(A)])$  and  $g(t) = R(1, \mathcal{D})f(t)$ , then we have

$$\begin{aligned} \|R(t)f\| &= \left\| \int_0^t T(s)f(-s) ds \right\| = \left\| \int_0^t T(s)(g(-s) - g'(-s)) ds \right\| \\ &\leq \left\| \int_0^t T(s)g(-s) ds \right\| + \left\| \int_0^t T(s)g'(-s) ds \right\| \\ &\leq \left\| \int_0^t T(s)g(-s) ds \right\| + \left\| \int_0^t T(s)Ag(-s) ds \right\| \end{aligned}$$

$$\begin{aligned}
& + \|T(t)g(-t)\| + \|g(0)\| \\
& \leq C \left( \sup_{s \in \mathbb{R}} \|g(s)\| + \sup_{s \in \mathbb{R}} \|Ag(s)\| \right) \\
& = C \|g\|_{BUC(\mathbb{R}, [D(A)])} = C \|R(1, \mathcal{D})f\|_{BUC(\mathbb{R}, [D(A)])} \\
& = C \|f\|_{BUC_{-1}(\mathbb{R}, [D(A)])}. \tag{2.25}
\end{aligned}$$

Here we used the fact that  $\int_0^t T(s)g'(-s)ds = \int_0^t AT(s)g(-s)ds - T(t)g(-t) + g(0)$ . Since  $BUC(\mathbb{R}, [D(A)])$  is dense in  $BUC_{-1}(\mathbb{R}, [D(A)])$ ,  $R(t)$  is uniformly bounded. By [Theorem 2.3](#), (2.24) has a unique bounded solution, and this concludes the proof.  $\square$

We complete this section with the following result, which is very helpful for studying properties of unbounded perturbed generators.

**THEOREM 2.10.** *For the operator matrix  $\mathcal{A} = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$  with  $D(\mathcal{A}) = D(A) \times D(B)$  assume that  $A$  is the generator of a  $C_0$ -semigroup on  $E$ ,  $B$  is the generator of a bounded  $C_0$ -group on  $F$  and  $C$  is  $B$ -bounded. Then  $\mathcal{A}$  is a generator of a  $C_0$ -semigroup on  $E \times F$  if and only if  $\mathcal{A}$  is of the form*

$$\mathcal{A} = \mathcal{D} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \mathcal{D}^{-1} + \mathcal{L} \tag{2.26}$$

with an isomorphism  $\mathcal{D}$  and a bounded operator  $\mathcal{L}$  in  $E \times F$ .

*Proof.* We have only to show the “only if” part. By the assumption we have  $\omega(S) = 0$ , where  $(S(t))_{t \in \mathbb{R}}$  is the group of  $-B$ . Without loss of generality, we assume  $\omega(T) < 0$ . Since  $\mathcal{A}$  is the generator of a  $C_0$ -semigroup on  $E \times F$ , by [6, Theorem 3.1], we have

$$V(t) := \int_0^t T(t-s)CS(-s)ds = \int_0^t T(s)CS(s-t)ds \tag{2.27}$$

is bounded for  $t \geq 0$ . Thus,

$$\begin{aligned}
\|R(t)f\| &= \left\| \int_0^t T(s)CS(s)f ds \right\| = \left\| \int_0^t T(s)CS(s-t)S(t)f ds \right\| \\
&\leq \|V(t)\| \cdot \|S(t)f\| \leq \|V(t)\| \cdot \|S(t)\| \cdot \|f\|.
\end{aligned} \tag{2.28}$$

Hence  $R(t)$  is bounded for every  $t \geq 0$ . By [Corollary 2.3](#), the equation  $AX - XB = C$  has a unique bounded solution  $X$ . Therefore,

$$\mathcal{A} = \begin{pmatrix} I & -X \\ 0 & I \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \begin{pmatrix} I & X \\ 0 & I \end{pmatrix}, \tag{2.29}$$

and the corollary is proved.  $\square$

### 3. Regularity of solutions of differential equations

In this section, we consider the differential equation on the line  $\mathbb{R}$

$$u'(t) = Au(t) + f(t), \quad t \in \mathbb{R}, \quad (3.1)$$

where  $A$  is the generator of a  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on  $E$  and  $f \in L^p(\mathbb{R}, E)$ . We say that the continuous function  $u(t)$  is a *mild solution* of (3.1), if

$$u(t) = T(t-s)u(s) + \int_s^t T(t-\tau)f(\tau)d\tau \quad (3.2)$$

for all  $t \geq s$ . It turns out that the existence and uniqueness of bounded mild solutions of (3.1) is closely related to the solvability of operator equations, as the following theorem shows.

**THEOREM 3.1.** *Let  $A$  be a generator of a  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  with  $\sigma(A) \cap i\mathbb{R} = \emptyset$ . Then the following are equivalent:*

- (a) *For each function  $f$  in  $L^p(\mathbb{R}, E)$  there exists a unique mild solution of (3.1), which is bounded.*
- (b) *There exists a unique bounded solution of the operator equation*

$$AX - X\mathcal{D} = -\delta_0, \quad (3.3)$$

where  $F = L^p(\mathbb{R}, E)$ ,  $\mathcal{D} : W^{p,1}(\mathbb{R}, E) \rightarrow F$ , and  $\delta_0 : W^{p,1}(\mathbb{R}, E) \rightarrow E$ .

*Proof.* (i)  $\Rightarrow$  (ii). Let  $C(\mathbb{R}, E)$  be the space of all bounded continuous functions over  $E$  and  $G : L^p(\mathbb{R}, E) \rightarrow C(\mathbb{R}, E)$  the operator defined by  $Gf = u$  where  $u$  is the unique mild solution of (3.1) corresponding to  $f$ . By standard argument, it is easy to see that  $G$  is a closed, and hence, a bounded operator. Define  $Xf = (Gf)(0)$ . Then  $X$  is a bounded operator from  $L^p(\mathbb{R}, E)$  to  $E$ .

Let now  $f \in W^{p,1}(\mathbb{R}, E)$ . By Corollary 2.7,  $u = Gf$  is a classical solution of (3.1), that is,

$$(Gf)'(t) = A(Gf)(t) + f(t). \quad (3.4)$$

Note that,  $(Gf)' = G(f')$ . Hence  $(Gf')(t) = A(Gf)(t) + f(t)$ . Taking  $t = 0$ , we have  $AXf - X\mathcal{D}f = -\delta_0 f$  for  $f \in \mathcal{D}$ , that is,  $X$  is a bounded solution of (3.3).

To show the uniqueness, we assume that  $X_0$  is a solution of equation  $AX - X\mathcal{D} = 0$ . Then for every  $f \in \mathcal{D}$  the function  $u \in \mathcal{M}$  defined by  $u(t) = XS(t)f$  is a classical solution of (3.1), since

$$u'(t) = X\mathcal{D}S(t)f = (AX + \delta_0)S(t)f = Au(t) + f(t) \quad (3.5)$$

for all  $t \in \mathbb{R}$ . Let now  $f \in \mathcal{M}$  and  $(f_k)_{k \in \mathbb{N}} \subseteq D(\mathcal{D})$  with  $\lim_k f_k = f$ . Then  $\lim_k u_k = \lim_k XS(\cdot)f_k = XS(\cdot)f$ . Hence, taking the limit on both sides of

$u_k = Gf_k$  as  $k \rightarrow \infty$  we get  $XS(\cdot)f = Gf$ , that is,  $u = XS(\cdot)f$  is a mild solution of (3.1). Assume now that  $X_1$  and  $X_2$  are two solutions of (3.3). Then, for every  $f \in \mathcal{M}$ ,  $u = (X_1 - X_2)S(\cdot)f$  is a mild solution (3.1). By the uniqueness of the mild solution we have  $u \equiv 0$ , which implies  $X_1 = X_2$ .

(ii) $\Rightarrow$ (i). We have shown above that, if  $X$  is a bounded solution of (3.3), then  $u(t) := XS(t)f$  is a mild solution of (3.1). It remains to be shown that this solution is unique. In order to do this, assume that  $v$  is a mild solution of the homogeneous equation  $u'(t) = Au(t)$ ,  $t \in \mathbb{R}$ . It is the well-known Tauberian theorem (cf. [11]) that the spectrum of the function  $f$ ,  $\text{sp}(f)$ , satisfies  $i\text{sp}(v) \subseteq \sigma(A)$ . By assumption,  $\sigma(A) \cap i\mathbb{R} = \emptyset$ , so that  $\text{sp}(v) = \emptyset$ , and so  $v \equiv 0$  (see [15, page 22]), and the theorem is proved.  $\square$

From [Corollary 2.5](#) and [Theorem 3.1](#) we obtain the following corollary.

**COROLLARY 3.2.** *If  $A$  is the generator of an exponentially dichotomic  $C_0$ -semigroup, then for every function  $f$  in  $L^p(\mathbb{R}, E)$ , (3.1) has a unique bounded mild solution.*

## References

- [1] W. Arendt, F. Räbiger, and A. Sourour, *Spectral properties of the operator equation  $AX + XB = Y$* , Quart. J. Math. Oxford Ser. (2) **45** (1994), no. 178, 133–149.
- [2] J. L. Daleckii and M. G. Krein, *Stability of Solutions of Differential Equations in Banach Space*, Translations of Mathematical Monographs, vol. 43, American Mathematical Society, Rhode Island, 1974. [MR 50#5126](#).
- [3] K. J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000. [MR 2000i:47075](#). [Zbl 0952.47036](#).
- [4] J. M. Freeman, *The tensor product of semigroups and the operator equation  $SX - XT = A$* , J. Math. Mech. **19** (1970), 819–828. [MR 43:7961](#). [Zbl 0206.44403](#).
- [5] G. Lumer and M. Rosenblum, *Linear operator equations*, Proc. Amer. Math. Soc. **10** (1959), 32–41. [MR 21:2927](#). [Zbl 0133.07903](#).
- [6] R. Nagel, *Towards a “matrix theory” for unbounded operator matrices*, Math. Z. **201** (1989), no. 1, 57–68. [MR 90c:47004](#). [Zbl 0672.47001](#).
- [7] R. Nagel and E. Sinestrari, *Inhomogeneous Volterra integrodifferential equations for Hille-Yosida operators*, Functional Analysis (Essen, 1991) (K. D. Bierstedt et al., eds.), Lecture Notes in Pure and Appl. Math., vol. 150, Dekker, New York, 1994, pp. 51–70. [MR 94i:34121](#). [Zbl 0790.45011](#).
- [8] T. L. Nguyen, *Higher differential equations and admissibility*, in preparation.
- [9] T. L. Nguyen and G. Nickel, *Time-dependent operator matrices and inhomogeneous Cauchy problems*, Rend. Circ. Mat. Palermo (2) **47** (1998), no. 1, 5–24. [MR 99f:34087](#). [Zbl 0916.47032](#).
- [10] V. Q. Phóng, *The operator equation  $AX - XB = C$  with unbounded operators  $A$  and  $B$  and related abstract Cauchy problems*, Math. Z. **208** (1991), no. 4, 567–588. [MR 93b:47035](#). [Zbl 0743.47022](#).

- [11] ———, *On the spectrum, complete trajectories, and asymptotic stability of linear semi-dynamical systems*, J. Differential Equations **105** (1993), no. 1, 30–45. [MR 94f:47049](#). [Zbl 0786.34068](#).
- [12] ———, *On the exponential stability and dichotomy of  $C_0$ -semigroups*, Studia Math. **132** (1999), no. 2, 141–149. [MR 2000j:47076](#). [Zbl 0926.47026](#).
- [13] V. Q. Phóng and E. Schüler, *The operator equation  $AX - XB = C$ , admissibility, and asymptotic behavior of differential equations*, J. Differential Equations **145** (1998), no. 2, 394–419. [MR 99h:34081](#). [Zbl 0918.34059](#).
- [14] J. Prüss, *On the spectrum of  $C_0$ -semigroups*, Trans. Amer. Math. Soc. **284** (1984), no. 2, 847–857. [MR 85f:47044](#).
- [15] ———, *Evolutionary Integral Equations and Applications*, Monographs in Mathematics, vol. 87, Birkhäuser Verlag, Basel, 1993. [MR 94h:45010](#).
- [16] M. Rosenblum, *On the operator equation  $BX - XA = Q$* , Duke Math. J. **23** (1956), 263–269. [MR 18:54d](#). [Zbl 0073.33003](#).
- [17] E. Schüler and V. Q. Phóng, *The operator equation  $AX - XD^2 = -\delta_0$  and second order differential equations in Banach spaces*, Semigroups of Operators: Theory and Applications (Newport Beach, CA, 1998), Progr. Nonlinear Differential Equations Appl., vol. 42, Birkhäuser, Basel, 2000, pp. 352–363. [MR 2001j:47020](#).

NGUYEN THANH LAN: DEPARTMENT OF MATHEMATICS, OHIO UNIVERSITY, ATHENS, OH 45701, USA

*E-mail address:* [tnguyen@math.ohio.edu](mailto:tnguyen@math.ohio.edu)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)