

*Research Article*

## **Positive Solutions of Boundary Value Problems for System of Nonlinear Fourth-Order Differential Equations**

Shengli Xie and Jiang Zhu

Received 23 March 2006; Revised 8 October 2006; Accepted 5 December 2006

Recommended by P. Joseph Mckenna

Some existence theorems of the positive solutions and the multiple positive solutions for singular and nonsingular systems of nonlinear fourth-order boundary value problems are proved by using topological degree theory and cone theory.

Copyright © 2007 S. Xie and J. Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### **1. Introduction and preliminary**

Fourth-order nonlinear differential equations have many applications such as balancing condition of an elastic beam which may be described by nonlinear fourth-order ordinary differential equations. Concerning the studies for singular and nonsingular case, one can refer to [1–10]. However, there are not many results on the system for nonlinear fourth-order differential equations. In this paper, by using topological degree theory and cone theory, we study the existence of the positive solutions and the multiple positive solutions for singular and nonsingular system of nonlinear fourth-order boundary value problems. Our conclusions and conditions are different from the ones used in [1–10] for single equations.

This paper is divided into three sections: in Section 2, we prove the existence of the positive solutions and the multiple positive solutions for systems of nonlinear fourth-order boundary value problems with nonlinear singular terms  $f_i(t, u)$  which may be singular at  $t = 0, t = 1$ . In Section 3, we prove some existence theorems of the positive solutions and the multiple positive solutions for nonsingular system of nonlinear fourth-order boundary value problems.

Let  $(E, \|\cdot\|)$  be a real Banach space and  $P \subset E$  a cone,  $B_\rho = \{u \in E : \|u\| < \rho\}$  ( $\rho > 0$ ).

## 2 Boundary Value Problems

LEMMA 1.1 [11]. *Assume that  $A : \bar{B}_\rho \cap P \rightarrow P$  is a completely continuous operator. If there exists  $x_0 \in P \setminus \{\theta\}$  such that*

$$x - Ax \neq \lambda x_0, \quad \forall \lambda \geq 0, x \in \partial B_\rho \cap P, \quad (1.1)$$

*then  $i(A, B_\rho \cap P, P) = 0$ .*

LEMMA 1.2 [12]. *Assume that  $A : \bar{B}_\rho \cap P \rightarrow P$  is a completely continuous operator and has no fixed point at  $\partial B_\rho \cap P$ .*

- (1) *If  $\|Au\| \leq \|u\|$  for any  $u \in \partial B_\rho \cap P$ , then  $i(A, B_\rho \cap P, P) = 1$ .*
- (2) *If  $\|Au\| \geq \|u\|$  for any  $u \in \partial B_\rho \cap P$ , then  $i(A, B_\rho \cap P, P) = 0$ .*

### 2. Singular case

We consider boundary value problems of singular system for nonlinear four order ordinary differential equations (SBVP)

$$\begin{aligned} x^{(4)} &= f_1(t, y), \quad t \in (0, 1), \\ x(0) = x(1) = x''(0) = x''(1) &= 0, \\ -y'' &= f_2(t, x), \quad t \in (0, 1), \\ y(0) = y(1) &= 0, \end{aligned} \quad (2.1)$$

where  $f_i \in C((0, 1) \times \mathbb{R}^+, \mathbb{R}^+)$  ( $i = 1, 2$ ),  $\mathbb{R}^+ = [0, +\infty)$ ,  $f_i(t, 0) \equiv 0$ ,  $f_i(t, u)$  are singular at  $t = 0$  and  $t = 1$ .  $(x, y) \in C^4(0, 1) \cap C^2[0, 1] \times C^2(0, 1) \cap C[0, 1]$  are a solution of SBVP (2.1) if  $(x, y)$  satisfies (2.1). Moreover, we call that  $(x, y)$  is a positive solution of SBVP (2.1) if  $x(t) > 0$ ,  $y(t) > 0$ ,  $t \in (0, 1)$ .

First, we list the following assumptions.

(H<sub>1</sub>) There exist  $q_i \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $p_i \in C((0, 1), [0, +\infty))$  such that  $f_i(t, u) \leq p_i(t)q_i(u)$  and

$$0 < \int_0^1 t(1-t)p_i(t)dt < +\infty \quad (i = 1, 2). \quad (2.2)$$

(H<sub>2</sub>) There exist  $\alpha \in (0, 1]$ ,  $0 < a < b < 1$  such that

$$\liminf_{u \rightarrow +\infty} \frac{f_1(t, u)}{u^\alpha} > 0, \quad \liminf_{u \rightarrow +\infty} \frac{f_2(t, u)}{u^{1/\alpha}} = +\infty \quad (2.3)$$

uniformly on  $t \in [a, b]$ .

(H<sub>3</sub>) There exists  $\beta \in (0, +\infty)$  such that

$$\limsup_{u \rightarrow 0^+} \frac{f_1(t, u)}{u^\beta} < +\infty, \quad \limsup_{u \rightarrow 0^+} \frac{f_2(t, u)}{u^{1/\beta}} = 0 \quad (2.4)$$

uniformly on  $t \in (0, 1)$ .

(H<sub>4</sub>) There exists  $\gamma \in (0, 1]$ ,  $0 < a < b < 1$  such that

$$\liminf_{u \rightarrow 0^+} \frac{f_1(t, u)}{u^\gamma} > 0, \quad \liminf_{u \rightarrow 0^+} \frac{f_2(t, u)}{u^{1/\gamma}} = +\infty \quad (2.5)$$

uniformly on  $t \in [a, b]$ .

(H<sub>5</sub>) There exists  $R > 0$  such that  $q_1[0, N] \int_0^1 t(1-t)p_1(t)dt < R$ , where  $N = q_2[0, R] \int_0^1 t(1-t)p_2(t)dt$ ,  $q_i[0, d] = \sup\{q_i(u) : u \in [0, d]\}$  ( $i = 1, 2$ ).

LEMMA 2.1 [13]. *Assume that  $p_i \in C((0, 1), [0, +\infty))$  ( $i = 1, 2$ ) satisfies (H<sub>1</sub>), then*

$$\lim_{t \rightarrow 0^+} t \int_t^1 (1-s)p_i(s)ds = \lim_{t \rightarrow 1^-} (1-t) \int_t^1 sp_i(s)ds = 0. \quad (2.6)$$

By (H<sub>1</sub>) and Lemma 2.1, we know that SBVP (2.1) is equivalent to the following system of nonlinear integral equations:

$$\begin{aligned} x(t) &= \int_0^1 G(t, s) \int_0^1 G(s, r) f_1(r, y(r)) dr ds, \\ y(t) &= \int_0^1 G(t, s) f_2(s, x(s)) ds, \end{aligned} \quad (2.7)$$

where

$$G(t, s) = \begin{cases} (1-t)s, & 0 \leq s \leq t \leq 1, \\ t(1-s), & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.8)$$

Clearly, (2.7) is equivalent to the following nonlinear integral equation:

$$x(t) = \int_0^1 G(t, s) \int_0^1 G(s, r) f_1 \left( r, \int_0^1 G(r, \tau) f_2(\tau, x(\tau)) d\tau \right) dr ds. \quad (2.9)$$

Let  $J = [0, 1]$ ,  $J_0 = [a, b] \subset (0, 1)$ ,  $\varepsilon_0 = a(1-b)$ ,  $E = C[0, 1]$ ,  $\|u\| = \max_{t \in J} |u(t)|$  for  $u \in E$ ,

$$K = \{u \in C[0, 1] : u(t) \geq 0, u(t) \geq t(1-t)\|u\|, t \in J\}. \quad (2.10)$$

It is easy to show that  $(E, \|\cdot\|)$  is a real Banach space,  $K$  is a cone in  $E$  and

$$\begin{aligned} G(t, s) &\geq \varepsilon_0, \quad \forall (t, s) \in J_0 \times J_0, \\ t(1-t)G(r, s) &\leq G(t, s) \leq G(s, s) = (1-s)s, \quad \forall t, s, r \in J. \end{aligned} \quad (2.11)$$

By virtue of (H<sub>1</sub>), we can define  $A : C[0, 1] \rightarrow C[0, 1]$  as follows:

$$(Ax)(t) = \int_0^1 G(t, s) \int_0^1 G(s, r) f_1(r, Tx(r)) dr ds, \quad (2.12)$$

## 4 Boundary Value Problems

where

$$(Tx)(t) = \int_0^1 G(t,s) f_2(s, x(s)) ds. \quad (2.13)$$

Then the positive solutions of SBVP (2.1) are equivalent to the positive fixed points of  $A$ .

LEMMA 2.2. *Let  $(H_1)$  hold, then  $A : K \rightarrow K$  is a completely continuous operator.*

*Proof.* Firstly, we show that  $T : K \rightarrow K$  is uniformly bounded continuous operator. For any  $x \in K$ , it follows from (2.13) that  $(Tx)(t) \geq 0$  and

$$(Tx)(t) \geq t(1-t) \int_0^1 G(r,s) f_2(s, x(s)) ds, \quad t, r \in J. \quad (2.14)$$

From (2.14), we get that  $(Tx)(t) \geq t(1-t) \|Tx\|$  for any  $t \in J$ . So  $T(K) \subset K$ .

Let  $D \subset K$  be a bounded set, we assume that  $\|x\| \leq d$  for any  $x \in D$ . Equation (2.13) and  $(H_1)$  imply that

$$\|Tx\| \leq q_2[0, d] \int_0^1 s(1-s) p_2(s) ds =: C_1, \quad (2.15)$$

from this we know that  $T(D)$  is a bounded set.

Next, we show that  $T : K \rightarrow K$  is a continuous operator. Let  $x_n, x_0 \in K$ ,  $\|x_n - x_0\| \rightarrow 0$  ( $n \rightarrow \infty$ ). Then  $\{x_n\}$  is a bounded set, we assume that  $\|x_n\| \leq d$  ( $n = 0, 1, 2, \dots$ ). By  $(H_1)$ , we have

$$\begin{aligned} f_2(t, x_n(t)) &\leq q_2[0, d] p_2(t), \quad t \in (0, 1), \quad n = 0, 1, 2, \dots, \\ |Tx_n(t) - Tx_0(t)| &\leq \int_0^1 s(1-s) |f_2(s, x_n(s)) - f_2(s, x_0(s))| ds, \quad t \in J. \end{aligned} \quad (2.16)$$

Now (2.16),  $(H_1)$ , and Lebesgue control convergent theorem yield

$$\|Tx_n - Tx_0\| \rightarrow 0 \quad (n \rightarrow \infty). \quad (2.17)$$

Thus  $T : K \rightarrow K$  is a continuous operator. By  $T \in C[K, K]$  and  $f_1 \in C((0, 1) \times \mathbb{R}^+, \mathbb{R}^+)$ , similarly we can show that  $A : K \rightarrow K$  is a uniformly bounded continuous operator.

We verify that  $A$  is equicontinuous on  $D$ . Since  $G(t, s)$  is uniformly continuous in  $J \times J$ , for any  $\varepsilon > 0$ ,  $0 \leq t_1 < t_2 \leq 1$ , there exists  $\delta = \delta(\varepsilon) > 0$  such that  $|t_1 - t_2| < \delta$  imply that  $|G(t_1, s) - G(t_2, s)| < \varepsilon$ ,  $s \in J$ . Then

$$\begin{aligned} |(Ax)(t_1) - (Ax)(t_2)| &\leq \int_0^1 |G(t_1, s) - G(t_2, s)| \int_0^1 G(s, r) f_1(r, (Tx)(r)) dr ds \\ &\leq q_1[0, C_1] \int_0^1 |G(t_1, s) - G(t_2, s)| ds \int_0^1 r(1-r) p_1(r) dr \\ &< \varepsilon q_1[0, C_1] \int_0^1 r(1-r) p_1(r) dr, \quad \forall x \in D. \end{aligned} \quad (2.18)$$

This implies that  $A(D)$  is equicontinuous. So  $A : K \rightarrow K$  is complete continuous. This completes the proof of Lemma 2.1.  $\square$

**THEOREM 2.3.** *Let  $(H_1)$ ,  $(H_2)$ , and  $(H_3)$  hold, then SBVP (2.1) has at least one positive solution.*

*Proof.* From Lemma 2.2, we know that  $A : K \rightarrow K$  is completely continuous. According to the first limit of  $(H_2)$ , there are  $\nu > 0$ ,  $M_1 > 0$  such that

$$f_1(t, u) \geq \nu u^\alpha, \quad \forall (t, u) \in [a, b] \times (M_1, +\infty). \quad (2.19)$$

Let  $R_1 \geq \max\{((b-a)\varepsilon_0^{(1+\alpha)/\alpha})^{-1}, ((\varepsilon_0^3(b-a)^2\nu/2) \max_{t \in J} \int_a^b G(t, s) ds)^{-1/\alpha}\}$ . By the second limit of  $(H_2)$ , there exists  $M_2 > 0$  such that

$$f_2(t, u) \geq R_1 u^{1/\alpha}, \quad \forall (t, u) \in [a, b] \times (M_2, +\infty). \quad (2.20)$$

Taking  $M \geq \max\{M_1, M_2\}$ ,  $\bar{R} = (M+1)\varepsilon_0^{-1}$ ,  $x_0(t) = \sin \pi t \in K \setminus \{\theta\}$ , we affirm that

$$x - Ax \neq \lambda x_0, \quad \forall \lambda \geq 0, x \in \partial B_{\bar{R}} \cap K. \quad (2.21)$$

In fact, if there are  $\lambda \geq 0$ ,  $x \in \partial B_{\bar{R}} \cap K$  such that  $x - Ax = \lambda x_0$ , then for  $t \in J$ , we have

$$x(t) \geq (Ax)(t) \geq \int_a^b G(t, s) \int_a^b G(s, r) f_1 \left( r, \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi \right) dr ds. \quad (2.22)$$

Owing to  $\alpha \in (0, 1]$  and  $x(t) \geq \varepsilon_0 \bar{R} > M$ ,  $t \in J_0$ , (2.20) implies that

$$\begin{aligned} \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi &\geq \int_a^b G(r, \xi) f_2(\xi, x(\xi)) d\xi \\ &\geq R_1 \int_a^b G(r, \xi) x^{1/\alpha}(\xi) d\xi \geq R_1 (\varepsilon_0 \bar{R})^{1/\alpha} \int_a^b G(r, \xi) d\xi \\ &\geq \bar{R} R_1 (b-a) \varepsilon_0^{1+1/\alpha} \geq \bar{R} > M, \quad r \in J_0. \end{aligned} \quad (2.23)$$

By using  $0 \leq G(t, s) \leq 1$ ,  $\alpha \in (0, 1]$  and Jensen inequality, it follows from (2.19)–(2.23) that

$$\begin{aligned} x(t) &\geq \nu \int_a^b G(t, s) \int_a^b G(s, r) \left( \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi \right)^\alpha dr ds \\ &\geq \varepsilon_0 \nu \int_a^b G(t, s) \int_a^b \left( \int_0^1 G^\alpha(r, \xi) f_2^\alpha(\xi, x(\xi)) d\xi \right) dr ds \\ &\geq \varepsilon_0 \nu \int_a^b G(t, s) \int_a^b \left( \int_a^b G(r, \xi) f_2^\alpha(\xi, x(\xi)) d\xi \right) dr ds \\ &\geq \varepsilon_0 \nu R_1^\alpha \int_a^b G(t, s) \int_a^b \int_a^b G(r, \xi) x(\xi) d\xi dr ds \\ &\geq \bar{R} \varepsilon_0^3 (b-a)^2 \nu R_1^\alpha \int_a^b G(t, s) ds, \quad t \in J. \end{aligned} \quad (2.24)$$

## 6 Boundary Value Problems

Thus

$$\bar{R} = \|x\| \geq \bar{R}\varepsilon_0^3(b-a)^2\nu R_1^\alpha \max_{t \in J} \int_a^b G(t,s)ds \geq 2\bar{R}. \quad (2.25)$$

This is a contradiction. By Lemma 1.1, we get

$$i(A, B_{\bar{R}} \cap K, K) = 0. \quad (2.26)$$

On the other hand, there exists  $\rho_1 \in (0, 1)$  according to the first limit of  $(H_3)$  such that

$$C_2 =: \sup \left\{ \frac{f_1(t, u)}{u^\beta} : (t, u) \in (0, 1) \times (0, \rho_1] \right\} < +\infty. \quad (2.27)$$

Taking  $\varepsilon_1 = \min\{\rho_1, (1/2C_2)^{1/\beta}\} > 0$ . By the second limit of  $(H_3)$ , there exists  $\rho_2 \in (0, 1)$  such that

$$f_2(t, u) \leq \varepsilon_1 u^{1/\beta}, \quad \forall (t, u) \in (0, 1) \times (0, \rho_2]. \quad (2.28)$$

Let  $\rho = \min\{\rho_1, \rho_2\}$ . Equations (2.27) and (2.28) imply that

$$\begin{aligned} \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi &\leq \varepsilon_1 \int_0^1 G(r, \xi) x(\xi)^{1/\beta} d\xi \leq \rho_1 \|x\|^{1/\beta} \\ &\leq \rho_1^{1+1/\beta} < \rho_1, \quad \forall x \in \bar{B}_\rho \cap K, r \in (0, 1), \\ (Ax)(t) &\leq C_2 \int_0^1 G(t, s) \left( \int_0^1 G(s, r) \left( \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi \right)^\beta dr \right) ds \\ &\leq C_2 \varepsilon_1^\beta \|x\| \leq \frac{1}{2} \|x\|, \quad \forall x \in \bar{B}_\rho \cap K, t \in [0, 1]. \end{aligned} \quad (2.29)$$

Then  $\|Ax\| \leq (1/2) \|x\| < \|x\|$  for any  $x \in \partial B_\rho \cap K$ . Lemma 1.2 yields

$$i(A, B_\rho \cap K, K) = 1. \quad (2.30)$$

Equations (2.26) and (2.30) imply that

$$i(A, (B_{\bar{R}} \setminus \bar{B}_\rho) \cap K, K) = i(A, B_{\bar{R}} \cap K, K) - i(A, B_\rho \cap K, K) = -1. \quad (2.31)$$

So  $A$  has at least one fixed point  $x \in (B_{\bar{R}} \setminus \bar{B}_\rho) \cap K$  which satisfies  $0 < \rho < \|x\| \leq \bar{R}$ . We know that  $x(t) > 0$ ,  $t \in (0, 1)$  by definition of  $K$ . This shows that SBVP (2.1) has at least one positive solution  $(x, y) \in C^4(0, 1) \cap C^2[0, 1] \times C^2(0, 1) \cap C[0, 1]$  by (2.7), and the solution  $(x, y)$  satisfies  $x(t) > 0$ ,  $y(t) > 0$  for any  $t \in (0, 1)$ . This completes the proof of Theorem 2.3.  $\square$

**THEOREM 2.4.** *Let  $(H_1)$ ,  $(H_4)$ , and  $(H_5)$  hold, then SBVP (2.1) has at least one positive solution.*

*Proof.* By the first limit of  $(H_4)$ , there exist  $\eta > 0$  and  $\delta_1 \in (0, R)$  such that

$$f_1(t, u) \geq \eta u^\gamma, \quad \forall (t, u) \in [a, b] \times [0, \delta_1]. \quad (2.32)$$

Let  $m \geq 2[\varepsilon_0^3(b-a)^2\eta \int_a^b G(1/2, s)ds]^{-1}$ , then according to the second limit of (H<sub>4</sub>), there exists  $\delta_2 \in (0, R)$  such that

$$f_2'(t, u) \geq mu, \quad \forall (t, u) \in [a, b] \times [0, \delta_2]. \quad (2.33)$$

Taking  $\delta = \min\{\delta_1, \delta_2\}$ . Since  $f_2(t, 0) \equiv 0$ ,  $f_2 \in C((0, 1) \times \mathbb{R}^+, \mathbb{R}^+)$ , there exists small enough  $\sigma \in (0, \delta)$  such that  $f_2(t, x) \leq \delta$  for any  $(t, x) \in (0, 1) \times [0, \sigma]$ . Then we have

$$\int_0^1 G(r, \tau) f_2(\tau, x(\tau)) d\tau \leq \delta, \quad \forall x \in \bar{B}_\sigma \cap K, r \in (0, 1). \quad (2.34)$$

By using Jensen inequality and  $0 < \gamma \leq 1$ , from (2.32)–(2.34) we can get that

$$\begin{aligned} (Ax) \left( \frac{1}{2} \right) &\geq \eta \int_a^b G\left(\frac{1}{2}, s\right) \int_a^b G(s, r) \left( \int_0^1 G(r, \tau) f_2(\tau, x(\tau)) d\tau \right)^\gamma dr ds \\ &\geq \varepsilon_0 \eta \int_a^b G\left(\frac{1}{2}, s\right) ds \int_a^b \left( \int_a^b G(r, \tau) f_2'(\tau, x(\tau)) d\tau \right) dr \\ &\geq \varepsilon_0 \eta m \int_a^b G\left(\frac{1}{2}, s\right) ds \int_a^b \int_a^b G(r, \tau) x(\tau) d\tau dr \\ &\geq \varepsilon_0^3 (b-a)^2 \eta m \|x\| \int_a^b G\left(\frac{1}{2}, s\right) ds \geq 2\|x\|, \quad \forall x \in \bar{B}_\sigma \cap K. \end{aligned} \quad (2.35)$$

From this we know that

$$\|Ax\| \geq 2\|x\| > \|x\|, \quad \forall x \in \partial B_\sigma \cap K. \quad (2.36)$$

Equation (2.36) and Lemma 1.2 imply that

$$i(A, B_\sigma \cap K, K) = 0. \quad (2.37)$$

On the other hand, for any  $x \in \partial B_R \cap K$ ,  $t \in [0, 1]$ , (H<sub>1</sub>) and (H<sub>5</sub>) imply that

$$\int_0^1 G(r, \tau) f_2(\tau, x(\tau)) d\tau \leq q_2[0, R] \int_0^1 \tau(1-\tau) p_2(\tau) d\tau = N, \quad (2.38)$$

$$\|Ax\| \leq q_1[0, N] \int_0^1 r(1-r) p_1(r) dr < R = \|x\|, \quad \forall x \in \partial B_R \cap K. \quad (2.39)$$

By (2.39) and Lemma 1.2, we obtain that

$$i(A, B_R \cap K, K) = 1. \quad (2.40)$$

Now, (2.37) and (2.40) imply that

$$i(A, (B_R \setminus \bar{B}_\sigma) \cap K, K) = i(A, B_R \cap K, K) - i(A, B_\sigma \cap K, K) = 1. \quad (2.41)$$

So  $A$  has at least one fixed point  $x \in (B_R \setminus \bar{B}_\sigma) \cap K$ , then SBVP (2.1) has at least one positive solution  $(x, y)$  which satisfies  $x(t) > 0$ ,  $y(t) > 0$  for any  $t \in (0, 1)$ . This completes the proof of Theorem 2.4.  $\square$

## 8 Boundary Value Problems

**THEOREM 2.5.** *Let  $(H_1)$ ,  $(H_2)$ ,  $(H_4)$ , and  $(H_5)$  hold, then SBVP (2.1) has at least two positive solutions.*

*Proof.* We take  $M > R > \sigma$  such that (2.26), (2.37), and (2.40) hold by the proof of Theorems 2.3 and 2.4. Then

$$\begin{aligned} i(A, (B_{\bar{R}} \setminus \bar{B}_R) \cap K, K) &= i(A, B_{\bar{R}} \cap K, K) - i(A, B_R \cap K, K) = -1, \\ i(A, (B_R \setminus \bar{B}_\sigma) \cap K, K) &= i(A, B_R \cap K, K) - i(A, B_\sigma \cap K, K) = 1. \end{aligned} \quad (2.42)$$

So  $A$  has at least two fixed points in  $(B_{\bar{R}} \setminus \bar{B}_R) \cap K$  and  $(B_R \setminus \bar{B}_\sigma) \cap K$ , then SBVP (2.1) has at least two positive solutions  $(x_i, y_i)$  and satisfies  $x_i(t) > 0$ ,  $y_i(t) > 0$  ( $i = 1, 2$ ) for any  $t \in (0, 1)$ . This completes the proof of Theorem 2.5.  $\square$

In the following, we give some applied examples.

*Example 2.6.* Let  $f_1(t, y) = y^2/t(1-t)$ ,  $f_2(t, x) = x^3/t(1-t)$ ,  $\alpha = \beta = 1/2$ . From Theorem 2.3, we know that SBVP (2.1) has at least one positive solution, here,  $f_1(t, y)$  and  $f_2(t, x)$  are superlinear on  $y, x$ , respectively.

*Example 2.7.* Let  $f_1(t, y) = y^{1/2}/t(1-t)$ ,  $f_2(t, x) = x^3/t(1-t)$ ,  $\alpha = \beta = 1/2$ . From Theorem 2.3, we know that SBVP (2.1) has at least one positive solution, here,  $f_1(t, y)$  and  $f_2(t, x)$  are sublinear and superlinear on  $y, x$ , respectively.

*Example 2.8.* Let  $f_1(t, y) = (y^2 + y^{1/2})/\sqrt{t(1-t)}$ ,  $f_2(t, x) = 4(x^3 + x^{1/2})/\pi\sqrt{t(1-t)}$ ,  $\alpha = \gamma = 1/2$ . It is easy to examine that conditions  $(H_1)$ ,  $(H_2)$ , and  $(H_4)$  of Theorem 2.5 are satisfied and  $\int_0^1 (dt/\sqrt{t(1-t)}) = \pi$ . In addition, taking  $R = 1$ , then  $q_2[0, 1] = \sup\{(x^3 + x^{1/2})/2 : x \in [0, 1]\} = 1$ ,  $N = (8/\pi)q_2[0, 1] \int_0^1 \sqrt{t(1-t)} dt = 1$ ,  $q_1[0, 1] = 2$ , where  $\int_0^1 \sqrt{t(1-t)} dt = \pi/8$ . Then  $q_1[0, 1] \int_0^1 \sqrt{t(1-t)} dt = \pi/4 < 1$ . Thus, the condition  $(H_5)$  of Theorem 2.5 is satisfied. From Theorem 2.5, we know that SBVP (2.1) has at least two positive solutions.

*Remark 2.9.* Balancing condition of a pair of elastic beams for fixed two ends may be described by boundary value problems for nonlinear fourth-order singular system (SBVP)

$$\begin{aligned} x^{(4)} &= f_1(t, -y''), \quad t \in (0, 1), \\ x(0) &= x(1) = x''(0) = x''(1) = 0, \\ y^{(4)} &= f_2(t, x), \quad t \in (0, 1), \\ y(0) &= y(1) = y''(0) = y''(1) = 0, \end{aligned} \quad (2.43)$$

where  $f_i \in C((0, 1) \times \mathbb{R}^+, \mathbb{R}^+)$  ( $i = 1, 2$ ),  $f_i(t, 0) \equiv 0$ ,  $f_i(t, u)$  are singular at  $t = 0$  and  $t = 1$ . Let  $-y''(t) = v(t)$ ,  $t \in [0, 1]$ . Then  $v(0) = v(1) = 0$ ,  $y(t) = \int_0^t G(t, s)v(s)ds$ , where  $G(t, s)$  is given by (2.8). SBVP (2.43) is changed into the form of SBVP (2.1)

$$\begin{aligned} x^{(4)} &= f_1(t, v), \quad t \in (0, 1), \\ x(0) &= x(1) = x''(0) = x''(1) = 0, \\ -v'' &= f_2(t, x), \quad t \in (0, 1), \\ v(0) &= v(1) = 0. \end{aligned} \quad (2.44)$$

Thus, from Theorems 2.3–2.5, we can get the existence of the positive solutions and multiple positive solutions of SBVP (2.43) under the conditions (H<sub>1</sub>)–(H<sub>5</sub>).

*Remark 2.10.* Balancing condition of a pair of bending elastic beams for fixed two ends may be described by boundary value problems for nonlinear fourth-order singular system (SBVP)

$$\begin{aligned} x^{(4)} &= f_1(t, -y''), \quad t \in (0, 1), \\ x(0) = x(1) &= x''(0) = x''(1) = 0, \\ y^{(4)} &= f_2(t, -x''), \quad t \in (0, 1), \\ y(0) = y(1) &= y''(0) = y''(1) = 0, \end{aligned} \tag{2.45}$$

where  $f_i \in C((0, 1) \times \mathbb{R}^+, \mathbb{R}^+)$  ( $i = 1, 2$ ),  $f_i(t, 0) \equiv 0$ ,  $f_i(t, u)$  are singular at  $t = 0$  and  $t = 1$ . Let  $-x''(t) = u(t)$ ,  $-y''(t) = v(t)$ ,  $t \in [0, 1]$ , then  $u(0) = u(1) = 0$ ,  $v(0) = v(1) = 0$  and the problem is equivalent to the following nonlinear integral equation system:

$$\begin{aligned} x(t) &= \int_0^1 G(t, s)u(s)ds, \\ y(t) &= \int_0^1 G(t, s)v(s)ds, \quad t \in [0, 1], \end{aligned} \tag{2.46}$$

where  $G(t, s)$  is given by (2.8). SBVP (2.45) is changed into the following boundary value problems for nonlinear second-order singular system:

$$\begin{aligned} -u'' &= f_1(t, v), \quad t \in (0, 1), \\ u(0) = u(1) &= 0, \\ -v'' &= f_2(t, u), \quad t \in (0, 1), \\ v(0) = v(1) &= 0. \end{aligned} \tag{2.47}$$

For SBVP (2.47), under conditions (H<sub>1</sub>)–(H<sub>5</sub>), by using the similar methods of our proof, we can show that SBVP (2.47) has the similar conclusions of Theorems 2.3–2.5.

### 3. Continuous case

We consider boundary value problems of system for nonlinear fourth-order ordinary differential equations (BVP)

$$\begin{aligned} x^{(4)} &= f_1(t, y), \quad t \in [0, 1], \\ x(0) = x(1) &= x''(0) = x''(1) = 0, \\ -y'' &= f_2(t, x), \quad t \in [0, 1], \\ y(0) = y(1) &= 0, \end{aligned} \tag{3.1}$$

where  $f_i \in C([0, 1] \times \mathbb{R}^+, \mathbb{R}^+)$ ,  $f_i(t, 0) \equiv 0$  ( $i = 1, 2$ ).  $(x, y) \in C^4[0, 1] \times C^2[0, 1]$  is a solution of BVP (3.1) if  $(x, y)$  satisfies (3.1). Moreover, we call that  $(x, y)$  is a positive solution of BVP (3.1) if  $x(t) > 0$ ,  $y(t) > 0$ ,  $t \in (0, 1)$ .

## 10 Boundary Value Problems

To prove our results, we list the following assumptions.

(Q<sub>1</sub>) There exists  $\tau \in (0, +\infty)$  such that

$$\limsup_{u \rightarrow +\infty} \frac{f_1(t, u)}{u^\tau} < +\infty, \quad \limsup_{u \rightarrow +\infty} \frac{f_2(t, u)}{u^{1/\tau}} = 0 \quad (3.2)$$

uniformly on  $t \in [0, 1]$ .

(Q<sub>2</sub>) There exists  $\beta \in (0, +\infty)$  such that

$$\limsup_{u \rightarrow 0^+} \frac{f_1(t, u)}{u^\beta} < +\infty, \quad \limsup_{u \rightarrow 0^+} \frac{f_2(t, u)}{u^{1/\beta}} = 0 \quad (3.3)$$

uniformly on  $t \in [0, 1]$ .

(Q<sub>3</sub>) There exist  $q_i \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $p_i \in C([0, 1], \mathbb{R}^+)$  such that  $f_i(t, u) \leq p_i(t)q_i(u)$  and there exists  $R > 0$  such that  $q_1[0, N] \int_0^1 t(1-t)p_1(t)dt < R$ , where  $N = q_2[0, R] \int_0^1 t(1-t)p_2(t)dt$ ,  $q_i[0, d] = \sup\{q_i(u) : u \in [0, d]\}$  ( $i = 1, 2$ ).

Obviously, for continuous case, the integral operator  $A : K \rightarrow K$  defined by (2.12) is complete continuous.

**THEOREM 3.1.** *Let (Q<sub>1</sub>) and (H<sub>4</sub>) hold. Then BVP (3.1) has at least one positive solution.*

*Proof.* We know that (2.37) holds by (H<sub>4</sub>). On the other hand, it follows from (Q<sub>1</sub>) that there are  $\omega > 0$ ,  $C_3 > 0$ ,  $C_4 > 0$  such that

$$\begin{aligned} f_1(t, u) &\leq \omega u^\tau + C_3, \quad \forall (t, u) \in [0, 1] \times \mathbb{R}^+, \\ f_2(t, u) &\leq \left(\frac{u}{2\omega}\right)^{1/\tau} + C_4, \quad \forall (t, u) \in [0, 1] \times \mathbb{R}^+. \end{aligned} \quad (3.4)$$

Noting that  $0 \leq G(t, s) \leq 1$ , (3.4) implies that

$$\begin{aligned} (Ax)(t) &\leq \int_0^1 G(t, s) \int_0^1 G(s, r) \left[ \omega \left( \int_0^1 G(r, \xi) f_2(\xi, x(\xi)) d\xi \right)^\tau + C_3 \right] dr ds \\ &\leq \omega \left( \int_0^1 \left[ \left( \frac{x(\xi)}{2\omega} \right)^{1/\tau} + C_4 \right] d\xi \right)^\tau + C_3 \leq \omega \left[ \left( \frac{\|x\|}{2\omega} \right)^{1/\tau} + C_4 \right]^\tau + C_3. \end{aligned} \quad (3.5)$$

By simple calculating, we get that

$$\lim_{\|x\| \rightarrow +\infty} \frac{\omega \left[ \left( \frac{\|x\|}{2\omega} \right)^{1/\tau} + C_4 \right]^\tau + C_3}{\|x\|} = \frac{1}{2}. \quad (3.6)$$

Then there exists a number  $G > 0$  such that  $\|x\| \geq G$  implies that

$$\omega \left[ \left( \frac{\|x\|}{2\omega} \right)^{1/\tau} + C_4 \right]^\tau + C_3 < \frac{3}{4} \|x\|. \quad (3.7)$$

Thus, we have

$$\|Ax\| < \|x\|, \quad \forall x \in \partial B_G \cap K. \quad (3.8)$$

It follows from (3.8) and Lemma 1.2 that

$$i(A, B_G \cap K, K) = 1. \quad (3.9)$$

Now, (2.37) and (3.9) imply that

$$i(A, (B_G \setminus \bar{B}_\sigma) \cap K, K) = i(A, B_G \cap K, K) - i(A, B_\sigma \cap K, K) = 1. \quad (3.10)$$

So  $A$  has at least one fixed point  $x \in (B_G \setminus \bar{B}_\sigma) \cap K$ , then BVP (3.1) has at least one positive solution  $(x, y)$  which satisfies  $x(t) > 0$ ,  $y(t) > 0$ ,  $t \in (0, 1)$ . This completes the proof of Theorem 3.1.  $\square$

Similar to the proof of Theorem 2.4, we can get the following theorems.

**THEOREM 3.2.** *Let  $(H_2)$  and  $(Q_2)$  hold. Then BVP (3.1) has at least one positive solution.*

**THEOREM 3.3.** *Let  $(H_4)$  and  $(Q_3)$  hold. Then BVP (3.1) has at least one positive solution.*

*Proof.* Similar to the proof of (2.37) and (2.40) in Section 2, we can prove that there exists  $\sigma \in (0, R)$  such that

$$i(A, B_\sigma \cap K, K) = 0, \quad i(A, B_R \cap K, K) = 1. \quad (3.11)$$

These imply that

$$i(A, (B_R \setminus \bar{B}_\sigma) \cap K, K) = i(A, B_R \cap K, K) - i(A, B_\sigma \cap K, K) = 1. \quad (3.12)$$

So  $A$  has at least one fixed point  $x \in (B_R \setminus \bar{B}_\sigma) \cap K$  and satisfies  $0 < \sigma < \|x\| \leq R$ . It follows from the definition of  $K$  that  $x(t) > 0$ ,  $t \in (0, 1)$ . This shows that BVP (3.1) has at least one positive solution  $(x, y)$  which satisfies  $x(t) > 0$ ,  $y(t) > 0$  for any  $t \in (0, 1)$ . This completes the proof of Theorem 3.3.  $\square$

From Theorems 3.2 and 3.3, we can get the following theorem.

**THEOREM 3.4.** *Let  $(H_2)$ ,  $(H_4)$ , and  $(Q_3)$  hold. Then BVP (3.1) has at least two positive solutions.*

In the following, we give some applications of Theorems 3.1–3.4.

*Example 3.5.* Let  $f_1(t, y) = y^{1/2}$ ,  $f_2(t, x) = x^{1/2}$ ,  $\tau = \gamma = 1/2$ . From Theorem 3.1, we know that BVP (3.1) has at least one positive solution, here,  $f_1(t, y)$  and  $f_2(t, x)$  are sublinear on  $y, x$ , respectively.

*Example 3.6.* Let  $f_1(t, y) = y^2$ ,  $f_2(t, x) = x^3$ ,  $\alpha = \beta = 1/2$ . From Theorem 3.2, we know that BVP (3.1) has at least one positive solution, here,  $f_1(t, y)$  and  $f_2(t, x)$  are superlinear on  $y, x$ , respectively.

*Example 3.7.* Let  $f_1(t, y) = y^{1/2}$ ,  $f_2(t, x) = x^3$ ,  $\alpha = \beta = 1/2$ . From Theorem 3.2, we know that BVP (3.1) has at least one positive solution, here,  $f_1(t, y)$  and  $f_2(t, x)$  are sublinear and superlinear on  $y, x$ , respectively.

## 12 Boundary Value Problems

*Example 3.8.* Let  $f_1(t, y) = (1/3)(y^2 + y^{1/2})$ ,  $f_2(t, x) = t(x^3 + x^{1/2})$ ,  $\alpha = \gamma = 1/2$ ,  $R = 1$ . From Theorem 3.4, we know that BVP (3.1) has at least two positive solutions.

*Remark 3.9.* From these examples we know that all conclusions in this paper are different from the ones in [1–10].

### Acknowledgments

The authors are grateful to the anonymous referee for his or her valuable comments. This work is supported by the Natural Science Foundation of the EDJP (05KGD110225), JSQLG, the National Natural Science Foundation 10671167, and EDAP2005KJ221, China.

### References

- [1] C. P. Gupta, “Existence and uniqueness theorems for the bending of an elastic beam equation,” *Applicable Analysis*, vol. 26, no. 4, pp. 289–304, 1988.
- [2] C. P. Gupta, “Existence and uniqueness results for the bending of an elastic beam equation at resonance,” *Journal of Mathematical Analysis and Applications*, vol. 135, no. 1, pp. 208–225, 1988.
- [3] R. P. Agarwal, “On fourth order boundary value problems arising in beam analysis,” *Differential Integral Equations*, vol. 2, no. 1, pp. 91–110, 1989.
- [4] D. O'Regan, “Solvability of some fourth (and higher) order singular boundary value problems,” *Journal of Mathematical Analysis and Applications*, vol. 161, no. 1, pp. 78–116, 1991.
- [5] Y. S. Yang, “Fourth-order two-point boundary value problems,” *Proceedings of the American Mathematical Society*, vol. 104, no. 1, pp. 175–180, 1988.
- [6] R. Ma and H. Wang, “On the existence of positive solutions of fourth-order ordinary differential equations,” *Applicable Analysis*, vol. 59, no. 1–4, pp. 225–231, 1995.
- [7] Z. L. Wei and Z. T. Zhang, “A necessary and sufficient condition for the existence of positive solutions of singular superlinear boundary value problems,” *Acta Mathematica Sinica*, vol. 48, no. 1, pp. 25–34, 2005 (Chinese).
- [8] Z. L. Wei, “Positive solutions to singular boundary value problems for a class of fourth-order sublinear differential equations,” *Acta Mathematica Sinica*, vol. 48, no. 4, pp. 727–738, 2005 (Chinese).
- [9] Y. X. Li, “Existence and multiplicity of positive solutions for fourth-order boundary value problems,” *Acta Mathematicae Applicatae Sinica*, vol. 26, no. 1, pp. 109–116, 2003 (Chinese).
- [10] Y. M. Zhou, “Positive solutions to fourth-order nonlinear eigenvalue problems,” *Journal of Systems Science and Mathematical Sciences*, vol. 24, no. 4, pp. 433–442, 2004 (Chinese).
- [11] D. J. Guo, *Nonlinear Functional Analysis*, Science and Technology Press, Jinan, Shandong, China, 1985.
- [12] D. J. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, vol. 5 of *Notes and Reports in Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1988.
- [13] J. G. Cheng, “Nonlinear singular boundary value problems,” *Acta Mathematicae Applicatae Sinica*, vol. 23, no. 1, pp. 122–129, 2000 (Chinese).

Shengli Xie: Department of Mathematics, Suzhou College, Suzhou 234000, China  
Email address: xieshengli200@sina.com

Jiang Zhu: School of Mathematics Science, Xuzhou Normal University, Xuzhou 221116, China  
Email address: jiangzhu@xznu.edu.cn

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)