

## THE DUAL SPACE OF $\beta(\Gamma)$

DENNIS NEMZER

Department of Mathematics  
California State University, Stanislaus  
Turlock, CA 95382

(Received August 17, 1995)

**ABSTRACT.** The space of Boehmians with  $\Delta$ -convergence is a complete topological vector space in which the topology is induced by an invariant metric. We show that the dual space of the space of periodic Boehmians can be identified with the class of trigonometric polynomials.

**KEY WORDS AND PHRASES:** Boehmian, dual space, generalized function.

**1991 AMS SUBJECT CLASSIFICATION CODES:** 44A40, 46A06, 46F99.

### 1. INTRODUCTION

In this note we will consider a class  $\beta$  of generalized functions known as Boehmians. The class of Boehmians is a generalization of regular operators, Schwartz distributions, and other spaces of generalized functions [1].

Mikusinski [2] has shown that  $\beta$  with  $\Delta$ -convergence is an  $F$ -space (i.e. a complete topological vector space, not necessarily convex, such that the topology is given by an invariant metric). Hence, a natural question arises; can we characterize  $\beta'$  the class of all continuous linear functionals on  $\beta$ ?

T. K. Boehme proved (unpublished) that the class  $\beta'(\mathcal{R})$  of all continuous linear functionals on  $\beta(\mathcal{R})$  (Boehmians on the real line) consists only of the trivial linear functional. That is, there are no nontrivial continuous linear functionals on  $\beta(\mathcal{R})$ . This is not the case for  $\beta'(\Gamma)$  (the dual space of the class of periodic Boehmians). Indeed, there are enough continuous linear functionals to separate points.

In this note, we will show that  $\beta'(\Gamma)$  can be identified with the class of all trigonometric polynomials.

Let  $\mathcal{A}$  denote the class of trigonometric polynomials. That is,  $p \in \mathcal{A}$  if for some  $m \in \mathbb{N}$ ,

$$p(x) = \sum_{n=-m}^m \alpha_n e^{inx}, \text{ where } \alpha_n \in \mathbb{C} \text{ for } n=0, \pm 1, \dots, \pm m.$$

**THEOREM 1.1.** For any  $T \in \beta'(\Gamma)$ , there exists a unique  $p(x) = \sum_{n=-m}^m \alpha_n e^{inx} \in \mathcal{A}$

(for some  $m \in \mathbb{N}$ ) such that

$$T(F) = \sum_{n=-m}^m \alpha_n \hat{F}(n), \text{ for all } F \in \beta(\Gamma). \quad (1.1)$$

Conversely, any  $p(x) = \sum_{n=-m}^m \alpha_n e^{inx} \in \mathcal{A}$  defines a continuous linear functional on  $\beta(\Gamma)$  via (1.1).

## 2. PRELIMINARIES

We use the same notation as in [3]. For other results concerning  $\beta(\Gamma)$  see [4,5]. In [3] the construction of  $\beta(\Gamma)$  differs from the construction in [4,5], but the space  $\beta(\Gamma)$  in both cases can be shown to be the same.

The collection of all continuous complex-valued functions on  $\mathcal{R}$  will be denoted by  $C(\mathcal{R})$ . The support of a continuous function  $f$ , denoted by  $\text{supp } f$ , is the complement of the largest open set on which  $f$  is zero.

The *convolution* of two continuous functions, where at least one has compact support, is given by  $(f*g)(x) = \int_{-\infty}^{\infty} f(x-t)g(t) dt$ .

A sequence of continuous nonnegative functions  $\{\delta_n\}$  will be called a *delta sequence* if (i)  $\int_{-\infty}^{\infty} \delta_n(x) dx = 1$  for  $n=1, 2, \dots$ , and (ii)  $\text{supp } \delta_n \subset (-\varepsilon_n, \varepsilon_n)$ , where  $\varepsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ .

A pair of sequences  $(f_n, \delta_n)$  is called a *quotient of sequences*, and denoted by  $f_n/\delta_n$ , if  $f_n \in C(\mathcal{R})$  ( $n=1, 2, \dots$ ),  $\{\delta_n\}$  is a delta sequence, and  $f_n * \delta_m = f_m * \delta_n$  for all  $m$  and  $n$ . Two quotients of sequences  $f_n/\delta_n$  and  $g_n/\sigma_n$  are equivalent if  $f_n * \sigma_m = g_m * \delta_n$  for all  $m$  and  $n$ . The equivalence classes are called *Boehmians*. The space of all Boehmians will be denoted by  $\beta(\mathcal{R})$ , and a typical element of  $\beta(\mathcal{R})$  will be written as  $F = f_n/\delta_n$ . By defining a natural addition and scalar multiplication on  $\beta(\mathcal{R})$ , i.e.  $f_n/\delta_n + g_n/\sigma_n = (f_n * \sigma_n + g_n * \delta_n)/(\delta_n * \sigma_n)$  and  $\alpha(f_n/\delta_n) = \alpha f_n/\delta_n$ , where  $\alpha$  is a complex number,  $\beta(\mathcal{R})$  becomes a vector space.

A sequence  $\{F_n\}$  of Boehmians is said to be  $\Delta$ -convergent to  $F$ , denoted by  $\Delta-\lim_{n \rightarrow \infty} F_n = F$ , if there exists a delta sequence  $\{\delta_n\}$  such that for each  $n$ ,  $(F_n - F) * \delta_n \in C(\mathcal{R})$  and  $(F_n - F) * \delta_n \rightarrow 0$  uniformly on compact sets as  $n \rightarrow \infty$ .

If either  $f$  is a periodic function of period  $2\pi$  or  $\text{supp } f \subset (-\pi, \pi)$ , the  $n^{\text{th}}$  *Fourier coefficient*  $\hat{f}(n)$  of  $f$  is defined as  $\hat{f}(n) = (2\pi)^{-1} \int_{-\pi}^{\pi} f(x) e^{-inx} dx$ , for  $n=0, \pm 1, \pm 2, \dots$ . By a simple calculation we see that  $(f * \delta)^{\hat{}}(n) = 2\pi \hat{f}(n) \delta(n)$  for all  $n$ .

Let  $\beta(\Gamma) = \{F \in \beta(\mathcal{R}) : F = \sum_{-\infty}^{\infty} a_n e^{inx}, \text{ for some sequence } \{a_n\} \text{ of complex numbers}\}$ .

That is  $F = \Delta-\lim_{n \rightarrow \infty} \sum_{k=-n}^n a_k e^{ikx}$ . For  $F \in \beta(\Gamma)$  such that  $F = \sum_{-\infty}^{\infty} a_n e^{inx}$ , the  $n^{\text{th}}$  *Fourier coefficient* of  $F$ , denoted by  $\hat{F}(n)$ , is  $\hat{F}(n) = a_n$ .

The proof of the next proposition is straightforward and hence is left to the reader.

**PROPOSITION 2.1.** Suppose that  $F_n \in \beta(\Gamma)$  for  $n=1, 2, \dots$ . If  $\Delta-\lim_{n \rightarrow \infty} F_n = F$ , then

$$\lim_{n \rightarrow \infty} \hat{F}_n(k) = \hat{F}(k) \text{ for each } k.$$

**PROPOSITION 2.2** [3, Lemma 3.5]. Let  $\{n_k\}$  be a subsequence of positive integers such that  $\sum_{k=1}^{\infty} 1/n_k < \infty$ . If  $\{a_n\}$  is any sequence of complex numbers such that  $a_n = 0$  for  $n \neq n_k$  ( $k=1, 2, \dots$ ),

then there is a Boehmian  $F \in \beta(\Gamma)$  such that  $\hat{F}(n) = a_n$  for all  $n$ .

The conclusion of Proposition 2.2 is also valid for a subsequence  $\{n_k\}$  of negative integers such that  $\sum_{k=1}^{\infty} 1/n_k > -\infty$ .

### 3. PROOF OF THE THEOREM

Suppose that  $T$  has the form of (1.1), then clearly  $T$  is linear. The continuity of  $T$  follows from Proposition 2.1.

For the other direction, suppose that  $T$  is any continuous linear functional on  $\beta(\Gamma)$ .

Let  $F = \sum_{-\infty}^{\infty} a_n e^{inx} \in \beta(\Gamma)$ . Then,  $T(F) = \sum_{-\infty}^{\infty} a_n T(e^{inx})$ .

Hence to finish the proof, it suffices to show that for only a finite number of  $n$ 's,  $T(e^{inx})$  does not vanish.

Suppose that this is not the case. Thus, there are infinitely many positive (negative)  $n$ 's such that  $T(e^{inx})$  does not vanish.

Now, pick a subsequence  $\{n_k\}$  from the above  $n$ 's such that  $n_k \geq 2^k$ , for  $k \in \mathbb{N}$ .

Let  $\alpha_{n_k} = (T(e^{in_k x}))^{-1}$ , for  $k \in \mathbb{N}$ .

By Proposition 2.2, there exists a  $G \in \beta(\Gamma)$  such that  $\hat{G}(n_k) = \alpha_{n_k}$  for  $k = 1, 2, \dots$  and zero otherwise. But this gives  $T(G) = \sum_{-\infty}^{\infty} \hat{G}(n) T(e^{inx}) = \infty$ .

Thus, for only a finite number of  $n$ 's,  $T(e^{inx})$  must not vanish and the proof is complete.

### 4. SOME CONCLUDING REMARKS

The dual space of  $\beta(\Gamma)$  can be identified with spaces other than the space of trigonometric polynomials, but these spaces are all isomorphic. For example, we could have identified  $\beta'(\Gamma)$  with the subspace of  $C^Z$  of finite sequences.

It is known that  $\beta(\Gamma)$  with  $\Delta$ -convergence is not a Banach space. Can the characterization of  $\beta'(\Gamma)$  in this paper be used to determine whether or not  $\beta(\Gamma)$  is a Fréchet space?

Does the strong convergence in  $\beta'(\Gamma)$  have a nice description in terms of some convergence of trigonometric polynomials?

Let  $\omega$  be a real-valued even function defined on the integers  $Z$  such that

$$0 = \omega(0) \leq \omega(n+m) \leq \omega(n) + \omega(m) \text{ for all } n, m \in Z \text{ and } \sum_{n=1}^{\infty} \frac{\omega(n)}{n^2} < \infty.$$

By using Theorems 3.1 and 3.2 in [5], we obtain the following interesting connection between weak convergence and  $\Delta$ -convergence in  $\beta(\Gamma)$ .

**COROLLARY 4.1.** Suppose that the set of positive integers is partitioned into two disjoint sets  $\{t_n\}$  and  $\{s_n\}$  such that  $\sum_{n=1}^{\infty} 1/t_n < \infty$ . Let  $\{F_n\}$  be a sequence of Boehmians such that the set

$\{e^{-\omega(s_k)} \hat{F}_n(\pm s_k)\}_{n=1}^{\infty}$  is uniformly bounded. Then a necessary and sufficient condition for the sequence

$\{F_n\}$  to be  $\Delta$ -convergent to zero is that  $\{F_n\}$  converges weakly to zero.

The space of periodic Beurling distributions  $P_{\omega}'$  can be identified with a subspace of Boehmians. By using Corollary 4.1 and the results in [6], one can show that the convergence structure in  $P_{\omega}'$  is stronger than the convergence  $P_{\omega}'$  inherits from  $\beta(\Gamma)$ .

**ACKNOWLEDGMENT.** The author would like to thank the referee for his/her helpful suggestions and comments.

#### REFERENCES

- [1] MIKUSIŃSKI, P., Boehmians and Generalized Functions, *Acta. Math. Hungarica* 51(1988), 271–281.
- [2] MIKUSIŃSKI, P., Convergence of Boehmians, *Japan. J. Math.* 9 (1983), 159–179.
- [3] NEMZER, D., The Product of a Function and a Boehmian, *Colloq. Math.* 66 (1993), 49–55.
- [4] NEMZER, D., Periodic Boehmians, *Internat. J. Math. & Math. Sci.* 12 (1989), 685–692.
- [5] NEMZER, D., Periodic Boehmians II, *Bull. Austral. Math. Soc.* 44 (1991), 271–278.
- [6] NEMZER, D., Convolution Quotients of Beurling Type, *Math. Japonica* 42 (1995), 1–8.

## Special Issue on Decision Support for Intermodal Transport

### Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

### Lead Guest Editor

**Gerrit K. Janssens**, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; [Gerrit.Janssens@uhasselt.be](mailto:Gerrit.Janssens@uhasselt.be)

### Guest Editor

**Cathy Macharis**, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; [Cathy.Macharis@vub.ac.be](mailto:Cathy.Macharis@vub.ac.be)