

RESEARCH NOTES  
ON THE CONVERGENCE OF FOURIER SERIES

GERALDO SOARES DE SOUZA

Department of Mathematics  
Auburn University  
Alabama 36849

(Received March 5, 1984)

ABSTRACT. We define the space  $B^p = \{f: (-\pi, \pi] \rightarrow \mathbb{R}, f(t) = \sum_{n=0}^{\infty} c_n b_n(t), \sum_{n=0}^{\infty} |c_n| < \infty\}$ .

Each  $b_n$  is a special  $p$ -atom, that is, a real valued function, defined on  $(-\pi, \pi]$ , which is either  $b(t) = 1/2\pi$  or  $b(t) = \frac{-1}{|I|^{1/p}} x_R(t) + \frac{1}{|I|^{1/p}} x_L(t)$ , where  $I$  is an interval in  $(-\pi, \pi]$ ,  $L$  is the left half of  $I$  and  $R$  is the right half.  $|I|$  denotes the length of  $I$  and  $x_E$  the characteristic function of  $E$ .  $B^p$  is endowed with the norm  $\|f\|_{B^p} = \inf \sum_{n=0}^{\infty} |c_n|$ , where the infimum is taken over all possible representations of  $f$ .  $B^p$  is a Banach space for  $1/2 < p < \infty$ .  $B^p$  is continuously contained in  $L^p$  for  $1 \leq p < \infty$ , but different. We have

THEOREM. Let  $1 < p < \infty$ . If  $f \in B^p$  then the maximal operator  $Tf(x) = \sup_n |S_n(f, x)|$  maps  $B^p$  into the Lorentz space  $L(p, 1)$  boundedly, where  $S_n(f, x)$  is the  $n^{\text{th}}$ -sum of the Fourier Series of  $f$ .

KEY WORDS AND PHRASES. *Maximal operator, Lorentz spaces and Fourier Series.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. Primary 42A20.

1. INTRODUCTION.

We define the space  $B^p = \{f: (-\pi, \pi] \rightarrow \mathbb{R}, f(t) = \sum_{n=0}^{\infty} c_n b_n(t), \sum_{n=0}^{\infty} |c_n| < \infty\}$ . Each  $b_n$  is a special  $p$ -atom that is, a real valued function  $b$ , defined on  $(-\pi, \pi]$ , which is either  $b(t) = 1/2\pi$  or  $b(t) = \frac{-1}{|I|^{1/p}} x_R(t) + \frac{1}{|I|^{1/p}} x_L(t)$ , where  $I$  is an interval in  $(-\pi, \pi]$ ,  $L$  is the left half of  $I$  and  $R$  is the right half.  $|I|$  denotes the length of  $I$  and  $x_E$  the characteristic function of  $E$ .  $B^p$  is endowed with the norm  $\|f\|_{B^p} = \inf \sum_{n=0}^{\infty} |c_n|$ , where the infimum is taken over all possible representations of  $f$ .  $B^p$  is a Banach space for  $1/2 < p < \infty$ .

These spaces were originally introduced by the author in [2]. Also see [3], [4], and [5].

In this note we are interested in  $p$  belonging to the interval  $(1, \infty)$ , that is,  $1 < p < \infty$ .

The Carleson-Hunt theorem on the almost everywhere convergence of a function  $f$  in  $L^p$  for  $1 < p < \infty$ , asserts that if  $f \in L^p$  then the Fourier Series of  $f$ , denoted by  $S(f, x)$ , converges to  $f$  almost everywhere. As is well known, this is a very powerful theorem in the theory of Fourier Analysis. However, many analysts agree that the proof of this theorem is too complicated to be accessible to the general audience, so that many researchers in the field have tried to give a suitable proof, but as far as I know nobody has done so. We refer the interested reader to [1] and [6].

In this note we present a proper subset of  $L^p$  for  $1 < p < \infty$ , namely the  $B^p$  functions, for which the proof of the convergence of Fourier Series is relatively simple.

The proof follows basically the idea of Carleson-Hunt, namely, we will prove that if  $f \in B^p$  then the operator defined by  $Tf(x) = \sup_n |S_n(f, x)|$  where  $S_n(f, x)$  is the  $n^{\text{th}}$ -sum of the Fourier series of  $f$ , is a bounded operator into the Lorentz space  $L(p, 1)$ .

Recall that a measurable function  $f$  belongs to the Lorentz space  $L(p, q)$  if  $\|f\|_{pq} = \left( \frac{1}{q} \int_0^\infty (f^*(t)t^{1/p})^q \frac{dt}{t} \right)^{1/q} < \infty$ , for  $0 < p \leq \infty$ ,  $0 < q \leq \infty$ , where  $f^*$  is the decreasing rearrangement of  $f$ , defined by  $f^*(t) = \text{Inf}\{y, |x: |f(x)| > y| \leq t\}$ , with  $|\cdot|$  being the Lebesgue measure. Observe that  $L(p, p)$  is the usual  $L^p$ -space. The space  $L(p, \infty)$  is also known as the weak  $L^p$ -space.

## 2. MAIN RESULTS.

The main result can be stated as follows.

**THEOREM A.** Let  $1 < p < \infty$ . If  $f \in B^p$  then the maximal operator defined by  $Tf(x) = \sup_n |S_n(f, x)|$  maps  $B^p$  into  $L(p, 1)$  boundedly, that is,  $\|Tf\|_{p1} \leq M \|f\|_{B^p}$

where  $M$  is a positive absolute constant, and  $S_n(f, x)$  being the  $n^{\text{th}}$ -sum of the Fourier series of  $f$ .

**PROOF.** First of all we notice that the operator  $T_a f = f^a$  where  $f^a(x) = f(x-a)$  maps  $B^p$  into  $B^p$  continuously, so that we just need to prove this result for  $f_h(t) = -\frac{1}{(2h)^{1/p}} x_{[-h, 0]}(t) + \frac{1}{(2h)^{1/p}} x_{[0, h]}(t)$ ,  $h > 0$  which will follow easily from

the estimate for  $g(t) = x_{[0, h]}(t)$ . In fact let  $S_n(g, x) = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t) D_n(x-t) dt$  where

$D_n(t) = \frac{\sin(n+1/2)t}{2 \sin t/2}$  is the Dirichlet Kernel. Therefore we have

$S_n(g, x) = \frac{1}{\pi} \int_{x-h}^x D_n(t) dt$ , we now use the elementary inequality  $|D_n(t)| \leq \frac{c}{|t|}, |t| \leq \pi$ ,

satisfied by the Dirichlet Kernels  $D_n$ , where  $c$  is an absolute constant. Thus

$$|S_n(g, x)| \leq \int_{x-h}^x |D_n(t)| dt \leq \int_{x-h}^x \frac{1}{|t|} dt \leq \frac{ch}{x-h} \leq \frac{2ch}{x}$$

for  $x > 2h$  and  $|S_n(g, x)| \leq \frac{2ch}{-x}$  for  $x < -2h$ .

I recall that  $\int_0^x D_n(t) dt$  is uniformly bounded in  $n$  and  $x$ , that is,

$$\left| \int_0^x D_n(t) dt \right| < A \text{ where } A \text{ is an absolute constant see [7], Volume 1, page 57.}$$

Consequently we have i)  $Tg(x) \leq A \forall x$  and ii)  $Tg(x) \leq \frac{2ch}{|x|}$  for  $|x| > 2h$ .

We now evaluate  $\|Tg\|_{p1}$  using the definition of  $L(p,1)$  norm. (See definition of  $L(p,1)$  given before) we have

$$\begin{aligned} \|Tg\|_{p1} &= \frac{1}{p} \int_0^\infty (Tg)^*(t) t^{\frac{1}{p}-1} dt = \frac{1}{p} \int_0^{2h} (Tg)^*(t) t^{\frac{1}{p}-1} dt + \frac{1}{p} \int_{2h}^\infty (Tg)^*(t) t^{\frac{1}{p}-1} dt \\ &\leq \frac{A}{p} \int_0^{2h} t^{\frac{1}{p}-1} dt + \frac{2ch}{p} \int_{2h}^\infty t^{\frac{1}{p}-2} dt = A(2h)^{\frac{1}{p}} + \frac{c}{p-1}(2h)^{\frac{1}{p}} \end{aligned}$$

for  $1 < p < \infty$ , therefore  $\|Tg\|_{p1} \leq M(2h)^{\frac{1}{p}}$  where  $M = A + \frac{c}{p-1}$ .

Now for  $f_h(t) = \frac{-1}{(2h)^{1/p}} x_{[-h,0]}(t) + \frac{1}{(2h)^{1/p}} x_{[0,h]}(t)$  we get  $\|Tf_h\|_{p1} \leq M$  and so

if  $f(t) = \sum_{n=0}^{\infty} c_n b_n(t)$  where the  $b_n$  are special  $p$ -atoms and  $\sum_{n=0}^{\infty} |c_n| < \infty$  we have

$$\|Tf\|_{p1} \leq M \sum_{n=0}^{\infty} |c_n| \text{ which implies } \|Tf\|_{p1} \leq M \|f\|_B^p. \text{ The proof is complete.}$$

COROLLARY. Let  $1 < p < \infty$ . If  $f \in B^p$  then  $S_n(f, x)$  converges to

$f(x)$  almost everywhere, where  $S_n(f, x)$  is the  $n^{\text{th}}$ -sum of the Fourier series of  $f$ .

PROOF. Let  $f$  in  $B^p$  for  $1 < p < \infty$ , then  $f(t) = \sum_{k=0}^{\infty} c_k b_k(t)$  where  $\sum_{k=0}^{\infty} |c_k| < \infty$

and the  $b_k$  are special  $p$ -atoms. Then we define the function  $w_f(x)$  by

$w_f(x) = \limsup_{n \rightarrow \infty} \left| S_n(f, x) - S_{\ell}(f, x) \right|$  so that,  $S_n(f, x)$  converges to  $f$  almost

$\begin{matrix} k > n \\ \ell \geq n \end{matrix}$

everywhere if and only if  $w_f(x) = 0$  almost everywhere.

Observe now that  $w_f(x) \leq 2Tf(x)$ . Therefore  $w_f \in L(p,1)$ . On the other hand we see that  $w_f(x) = w_{f-f_n}(x)$  where  $f_n \rightarrow f$  in the  $B^p$ -norm, namely take  $f_n(t) =$

$\sum_{k=0}^n c_k b_k(t)$  a finite linear combination of special  $p$ -atoms  $b_k$ , then

$w_f(x) = w_{f-f_n}(x) \leq 2T(f-f_n)(x)$  and consequently theorem A, that is, the boundedness

of  $T$ , implies that  $\|w_f\|_{p1} \leq 2\|T(f-f_n)\|_{p1} \leq 2M\|f-f_n\|_{B^p}$ , so that as  $\|f - f_n\|_{B^p} \rightarrow 0$

for  $n \rightarrow \infty$ , we get  $\|w_f\|_{p1} = 0$ , thus  $w_f(x) = 0$  almost everywhere, which implies

$S_n(f, x) \rightarrow f(x)$  almost everywhere. The proof is complete.

We would like to point out that other direct proofs of the almost everywhere convergence for functions in  $B^p$  are also available. However we prefer this one because it is a consequence of the boundedness of the maximal operator and this boundedness could be useful in other contexts, as for example in interpolation of operators.

Another way of proving the almost everywhere convergence is by observing that if  $f$  is in  $B^p$  then

$$\int_0^{2\pi} \int_0^{2\pi} \left| \frac{f(x) - f(y)}{x - y} \right| dx dy < \infty.$$

The proof of this assertion is elementary and will be left to the interested reader, however we point out that we just need to prove it for functions of the type

$$f(x) = \frac{1}{h^{1/p}} x^p (x), \quad [0, h]$$

A consequence of the boundedness of the above integral is that any  $f$  in  $B^p$  satisfies Dini's condition and therefore the almost everywhere convergence is readily established.

One of the important features of the spaces  $B^p$  for  $1 \leq p < \infty$  is that it can be identified with the space of analytic functions  $F$  on the disk

$$D = \{z \in C: |z| < 1\} \text{ satisfying} \\ \int_0^1 \int_0^{2\pi} |F'(re^{i\theta})| (1-r)^{1/p-1} dr d\theta < \infty$$

where the dash means derivative.

ACKNOWLEDGEMENT: This paper is dedicated to Jonio Lemos, my teacher. We would like to thank R. A. Zalik who read the manuscript and made useful comments. We also wish to thank the referee for his helpful suggestions.

#### REFERENCES

1. Carleson, L., On convergence and growth of partial sums of Fourier series, Acta Math. 116(1966) 135-157.
2. De Souza, G. S., Spaces formed by special atoms, Ph.D. dissertation, SUNY at Albany, New York, 1980.
3. De Souza, G. S., Spaces formed by special atoms I, to appear in the Rocky Mountain Journal of Mathematics.
4. De Souza, G. S., Spaces formed by special atoms II, Functional Analysis, Holomorphy and Approximations Theory II, G. I. Zapata (Ed.), Elsevier Science Publishers B.V. North-Holland 1984.
5. De Souza, G. S., Two Theorems on Interpolation of Operators, Journal of Functional Analysis 46, 149-157(1982).
6. Hunt, R. A., On the Convergence of Fourier Series, Proc. Conference Southern Illinois University, Edwardsville, Illinois, 1967.
7. Zygmund, A., Trigonometric Series, Cambridge University Press, London and New York, 1959.

## Special Issue on Space Dynamics

### Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

#### Lead Guest Editor

**Antonio F. Bertachini A. Prado**, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; [prado@dem.inpe.br](mailto:prado@dem.inpe.br)

#### Guest Editors

**Maria Cecilia Zanardi**, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; [cecilia@feg.unesp.br](mailto:cecilia@feg.unesp.br)

**Tadashi Yokoyama**, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; [tadashi@rc.unesp.br](mailto:tadashi@rc.unesp.br)

**Silvia Maria Giuliatti Winter**, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; [silvia@feg.unesp.br](mailto:silvia@feg.unesp.br)